

July 2017

Studies on the Feculae of Selected Michigan Acrididae (Orthoptera)

Michael Tyrkus
Wayne State University

S.K. Gangwere
Wayne State University

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

Tyrkus, Michael and Gangwere, S.K. 2017. "Studies on the Feculae of Selected Michigan Acrididae (Orthoptera)," *The Great Lakes Entomologist*, vol 3 (4)

DOI: <https://doi.org/10.22543/0090-0222.1130>

Available at: <https://scholar.valpo.edu/tgle/vol3/iss4/5>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

STUDIES ON THE FECULAE OF SELECTED MICHIGAN ACRIDIDAE (ORTHOPTERA)¹

Michael Tyrkus and S. K. Gangwere

Department of Biology,
Wayne State University, Detroit, Michigan 48202

INTRODUCTION

Feculae have been used for some years in the study of insect food-habits and biology. Among the past authors who have used them are Frost (1928), Hodson and Brooks (1956), and Morris (1942), who discussed the use of feculae and frass in the identification of insect species in general; Boldyrev (1928), who figured the feculae of the bradyporine katydid *Bradyporus multituberculatus*; Brown (1937), who described in detail fecular microstructure in the spine-breasted grasshopper *Melanoplus bivittatus*; Weiss and Boyd (1950, 1952), who figured the feculae of representatives of selected insect orders; Gangwere (1962), who described the various factors responsible for shaping and coloring feculae, and with Morales (Gangwere and Morales, 1964) erected the first formal classification of orthopteroid feculae, and later (Gangwere, 1969) discussed the use of feculae of museum specimens as a shortcut method for determining food-habit; Scott (1964), who constructed a pictorial key to the droppings of cockroaches; Simeone (1964), who described the frass of posting beetles; Bhatia and Chandra (1967), who described the effect of food plants on the color, structure, and size of feculae in the locust *Schistocerca gregaria*; and Sardesai (1969) who reported on size variation in feculae of larval Lepidoptera.

With the partial exception of Gangwere and Morales (1964) and Bhatia and Chandra (1967), investigators have paid scant attention to variations in fecular structure within given species. The present study fills, in part, this gap in our knowledge. It describes those differences in fecular size and structure in selected Michigan acridids that relate to sex of defecator and type of food. It also attempts a statistical measurement of such differences.

MATERIALS AND METHODS

During the field seasons of 1964 and 1965, individuals of several species of Acrididae were collected at the Edwin S. George Reserve, Livingston County, Michigan, and, during the summer of 1968, at the Proud Lake and the Pontiac Lake Recreation Areas, Oakland County, and the Stony Creek Metropolitan Park, Macomb County, Michigan. The captured insects were maintained both in the field and in the laboratory using the rearing cages described in Gangwere (1960).

All animals from the E. S. George Reserve were caged in the field. They were isolated in individual mantid-type cages, as described by Gangwere (1960). The cages were placed on a table within a box-like wire window screening enclosure approximately 4 ft. x 4 ft. x 3 ft., and the table, in turn, was left uncovered in the field. Inasmuch as the cages were exposed to the out-of-door conditions, all testing was necessarily restricted to periods of good weather. Preparatory to a given test, each individual animal was starved for a period of 24 hours (emptying the gut), and then was allowed to feed on a single, known species of plant presented in a small Erlenmeyer flask. No other food was given, though each insect had access to water from a cotton-stoppered vial. Both animals and feculae were needed for study and measurement; thus, at the end of the feeding period, the insects were killed and preserved in 80% alcohol, and their feculae were harvested from the cage floors.

The animals from all stations other than the E. S. George Reserve were treated in a similar manner, except that all tests upon them were carried out in the laboratory and in absence of the protective screen enclosure.

¹Contribution No. 246 from the Department of Biology, Wayne State University, Detroit, Michigan 48202. Based in part on a 1969 thesis submitted by Tyrkus in partial fulfillment of the requirements for the degree of Master of Science, Wayne State University, and in part research by Gangwere.

The length of all feculae was measured in the laboratory under a binocular compound microscope using an ocular micrometer. The length of the animals was taken with the method of Gangwere and Morales (1964); it was based on the distance between the anteriormost extension of the head capsule and the apex of the hind femur (with the latter placed so that its long axis is parallel to the body). This measurement was accomplished under a binocular dissecting microscope by placing the animal in question against a rule, and was standardized by use of an ocular micrometer.

The species investigated include the following:

Acrididae: Oedipodinae (Band-Winged Grasshoppers)

Arphia p. pseudonietana (Thomas)

Spharagemon b. bolli Scudder

Spharagemon collare (Scudder)

Acrididae: Catantopinae (Spine-Breasted Grasshoppers)

Melanoplus confusus Scudder

Melanoplus s. sanguinipes (Fabricius)

RESULTS

The results of the study are presented entirely in tabular form. They include Table 1 (Studies on the Feculae of Selected Acrididae: Basic Data), Table 2A (Size Variation in Defecating Animals and their Respective Feculae), Table 2B (Size Variation in Defecating Animals and their Feculae: By Fecular Type), and Table 3 (One-Way Analysis of Variance for Variable Ratios Between Various Populations and Groups).

Table 1. Studies on the feculae of selected Acrididae: basic data.

Plants ingested	Feeders ²	Sex	Ave. Length Feculae (mm)	Ave. Length Defecators (mm)	Ratio of Lengths	Types of Feculae ³
Equisetaceae						
<i>Equisetum arvense</i>	MC	F	5.10	22	1:4.31	1B
Moss sp.	SC	M	3.96	20	1:5.05	1C
Gramineae						
<i>Aristida purpurascens</i>	MC	F	4.65	21	1:4.52	1C
" "	MS	F	5.34	22	1:4.10	1B
" "	AP	F	7.56	23	1:3.04	1B
" "	SB	F	6.14	28	1:4.56	1B
" "	SC	M	4.39	22	1:5.02	1B
<i>Cenchrus sp.</i>	AP	M	3.83	21	1:5.49	1B
<i>Danthonia spicata</i>	MC	F	4.50	20	1:4.44	1B
" "	SC	F	6.24	22	1:3.53	1B
<i>Eragrostis spectabilis</i>	SB	F	4.26	—	—	1B
<i>Leptoloma cognatum</i>	MS	F	4.97	20	1:4.01	BC
" "	AP	F	7.16	22	1:3.07	1B
" "	SC	M	4:12	19	1:4.61	BC
<i>Panicum depauperatum</i>	SC	M	2.67	19	1:7.10	1B
" "	SC	F	5.21	24	1:4.61	1B
<i>Panicum oligosenthes</i>	MC	F	4.01	19	1:4.01	1B
" "	MS	M	4.75	17	1:3.58	BC
" "	AP	M	4.21	21	1:4.99	1B
" "	SB	F	3.17	26	1:8.20	1B
" "	SC	M	4.19	19	1:4.54	1B

Table 1 (Continued)

<i>Poa compressa</i>	MC	F	4.56	21	1:4.61	1B
" "	MS	M	4.42	16	1:3.62	1B
" "	AP	M	5.25	23	1:4.38	1B
" "	AP	F	5.25	25	1:4.76	1B
" "	SB	F	5.88	26	1:4.42	1B
" "	SC	F	4.96	22	1:4.43	1B
<i>Poa pratensis</i>	MS	F	5.34	22	1:4.10	1B
" "	AP	M	---	21	----	1B
" "	SB	M	3.22	-	----	1B
" "	SC	M	3.47	21	1:6.05	1B
" "	SC	M	4.42	20	1:4.52	1B
<i>Setaria glauca</i>	AP	F	4.70	24	1:5.11	1B
" "	SC	F	5.31	25	1:4.71	1B
Cyperaceae						
<i>Carex muhlenbergii</i>	MC	M	2.08	18	1:8.65	BC
<i>Cyperus filiculmis</i>	MC	M	3.28	16	1:4.88	1B
" "	MS	M	3.41	20	1:5.86	BC
" "	AP	M	7.37	22	1:2.98	1B
" "	SB	F	5.03	-	----	1B
" "	SC	M	4.16	19	1:4.57	1C
Cupressaceae						
<i>Juniperus sp.</i>	MC	F	4.16	22	1:5.29	1B
Polygonaceae						
<i>Rumex acetosella</i>	MC	F	2.97	22	1:7.41	1B
" "	MS	M	3.55	18	1:5.08	1C
" "	AP	M	3.07	22	1:7.17	BC
" "	SB	F	6.14	29	1:4.72	1C
" "	SC	F	4.21	23	1:5.46	1C
Cruciferae						
<i>Lepidium virginicum</i>	MC	F	3.44	19	1:5.52	1C
Rosaceae						
<i>Potentilla intermedia</i>	MC	F	3.32	19	1:5.72	1B
" "	SC	M	3.73	20	1:5.36	1C
" "	SC	M	3.53	19	1:5.38	1C
<i>Rubus flagellaris</i>	MC	M	2.28	17	1:8.17	1C
" "	MC	M	2.50	17	1:6.80	1C
" "	MS	F	3.37	-	----	1C
" "	AP	M	3.92	22	1:5.61	1C
" "	SC	M	3.00	20	1:6.67	1C
Juglandaceae						
<i>Carya ovalis</i>	MS	F	2.89	20	1:6.92	1C
" "	SC	M	3.07	19	1:6.19	1C
<i>Carya sp.</i>	MC	F	3.12	19	1:6.09	1B
" "	MC	M	2.35	-	----	1C
Fabaceae						
<i>Desmodium illinoense</i>	MC	M	2.48	17	1:6.88	1C
" "	AP	M	3.47	21	1:6.05	BC
" "	AP	F	6.31	28	1:4.43	BC
<i>Desmodium sessilifolium</i>	AP	M	3.41	21	1:6.15	1C
<i>Lespedeza capitata</i>	MC	F	4.28	19	1:4.49	1C
" "	MS	F	3.47	19	1:5.47	1C
" "	AP	F	9.56	26	1:2.72	1C

Table 1 (Continued)

<i>Lespedeza capitata</i>	SB	F	5.43	27	1:4.97	1B
" "	SC	F	4.55	24	1:5.27	1C
<i>Lespedeza hirta</i>	MC	M	2.72	17	1:6.25	1C
" "	AP	F	3.66	24	1:6.56	1C
" "	SC	F	5.15	25	1:4.85	--
Oxalidaceae						
<i>Oxalis stricta</i>	MC	F	3.66	22	1:6.01	1B
" "	MS	F	3.56	20	1:5.62	1C
" "	AP	M	2.18	22	1:10.09	1C
" "	SC	M	4.11	20	1:4.87	1C
Fagaceae						
<i>Quercus rubra</i>	SC	F	4.13	22	1:5.33	1C
<i>Quercus</i> sp.	MC	M	2.65	21	1:7.92	1C
Euphorbiaceae						
<i>Euphorbia corollata</i>	AP	F	3.33	25	1:7.51	BC
" "	SB	F	2.89	30	1:10.38	1C
" "	SC	F	5.44	24	1:4.41	BC
Hypericaceae						
<i>Hypericum perforatum</i>	MC	M	2.72	17	1:6.25	1C
" "	MC	F	4.21	21	1:4.99	1C
" "	MS	M	3.27	20	1:6.11	1C
" "	SC	F	3.47	23	1:6.63	1C
" "	SC	F	5.74	23	1:4.01	1C
Cistaceae						
<i>Helianthemum canadense</i>	MC	M	2.45	17	1:6.94	1C
<i>Lechea villosa</i>	MS	F	3.68	22	1:5.99	1C
" "	AP	F	3.96	26	1:6.57	1C
" "	SC	M	2.73	21	1:7.69	1C
Umbelliferae						
<i>Daucus carota</i>	MC	F	3.94	23	1:5.85	1C
Asclepiadaceae						
<i>Asclepias syriaca</i>	MC	F	3.57	19	1:5.32	1C
" "	MC	F	3.31	20	1:6.05	1C
" "	MS	F	2.90	20	1:6.89	1C
" "	SB	F(nymph)	6.11	22	1:3.60	1C
" "	SC	F	2.57	20	1:7.79	1C
Verbenaceae						
<i>Verbena</i> sp.	MC	F	4.19	20	1:4.77	1C
Labiatae						
<i>Leonurus cardiaca</i>	MC	F	4.06	21	1:5.17	1C
<i>Monarda fistulosa</i>	MS	F	2.97	22	1:7.41	1C
" "	AP	M	3.76	21	1:5.58	1C
" "	SC	F	4.90	24	1:4.90	1C
Solanaceae						
<i>Physalis heterophylla</i>	MC	F	3.86	20	1:5.18	1C
" "	MS	F	4.03	21	1:5.21	1C
" "	SC	M	2.67	18	1:6.74	BC
Plantaginaceae						
<i>Plantago aristata</i>	MC	F	2.87	21	1:7.32	1C
" "	MS	M	3.17	18	1:5.68	1C
" "	SC	M	4.16	20	1:4.81	1C
" "	SC	M	4.26	19	1:4.46	1C

Table 1 (Continued)

<i>Plantago rugelii</i>	SC	M	3.08	21	1:6.82	1C
Compositae						
<i>Achillea millefolium</i>	AP	F	6.14	25	1:4.07	1C
" "	SC	M	4.36	20	1:4.59	1C
<i>Ambrosia artemisiifolia</i>	MC	F	2.99	21	1:7.02	1C
" "	AP	F	3.91	24	1:6.14	1C
" "	SB	M	3.80	22	1:5.79	1C
" "	SC	M	2.77	19	1:6.86	1C
<i>Antennaria fallax</i>	MC	M	3.17	19	1:6.00	1C
<i>Antennaria neglecta</i>	MC	M	2.63	18	1:6.85	1C
" "	MC	F	3.47	20	1:5.76	1C
" "	MS	M	2.89	19	1:6.59	1C
" "	AP	M	3.96	21	1:5.33	1C
" "	SC	M	3.81	20	1:5.25	BC
" "	SC	M	3.62	20	1:5.52	BC
<i>Arctium minus</i>	MC	F	4.60	20	1:4.35	1C
<i>Chrysanthemum leucanthemum</i>	MC	M	3.12	18	1:5.77	1C
<i>Erigeron strigosus</i>	MC	F	4.36	25	1:5.71	1C
" "	MS	F	4.21	22	1:5.22	1C
" "	SB	F	4.72	—	—	1C
" "	SC	F	3.46	24	1:6.94	1C
<i>Gnaphalium obtusifolium</i>	AP	F	4.06	25	1:6.15	1C
" "	SC	F	2.09	24	1:11.48	1C
<i>Hieracium gronovii</i>	MC	F	3.83	21	1:5.48	1C
" "	SB	F	5.45	25	1:4.59	1C
" "	SC	F	3.40	25	1:7.35	1C
<i>Hieracium longipilum</i>	MC	F	4.85	22	1:4.54	1C
" "	MC	M	3.25	17	1:5.23	1C
" "	MS	M	4.21	19	1:4.51	1C
" "	MS	F	4.23	15	1:3.54	1C
" "	AP	F	6.47	26	1:4.02	BC
" "	AP	M	3.71	22	1:5.93	1C
" "	SB	F	6.19	28	1:4.52	1C
" "	SC	M	4.31	19	1:4.41	BC
" "	SC	M	4.16	19	1:4.57	BC
<i>Hieracium sp.</i>	MS	F	3.56	21	1:5.89	1C
<i>Lactuca sp.</i>	MC	F	4.39	20	1:4.55	1C
<i>Liatris aspera</i>	MC	M	2.70	17	1:6.28	1C
" "	MC	M	2.92	18	1:6.16	1C
" "	MS	M	3.31	17	1:5.13	1C
" "	SB	M	2.81	21	1:7.47	1C
" "	SC	M	5.34	19	1:3.55	1C
<i>Solidago juncea</i>	MC	F	3.09	21	1:6.79	1C
" "	MC	F	4.61	23	1:4.99	1C
" "	MC	F	3.04	19	1:6.25	1C
" "	MS	M	3.76	19	1:5.05	1C
<i>Solidago nemoralis</i>	MC	F	3.52	21	1:5.97	1C
" "	MS	M	2.77	18	1:6.85	1C
" "	SB	F	4.46	31	1:6.95	1C
" "	SB	M	2.65	20	1:7.55	1C
<i>Solidago rigida</i>	MS	F	3.63	20	1:5.50	1C
" "	AP	M	3.78	22	1:5.82	BC
" "	SB	F	3.60	—	—	1C

Table 1 (Continued)

<i>Taraxacum officinale</i>	AP	M	—	23	---	1C
<i>Tragopogon pratensis</i>	MC	F	4.21	21	1:4.99	1C
”	MC	F	5.77	21	1:3.64	1C
”	MS	F	3.36	20	1:6.00	1C
”	SC	F	4.65	23	1:4.95	1C
<i>Verbascum thapsus</i>	SC	M	3.10	18	1:5.81	1C

²Feeding species are denoted by the following code: MC = *Melanoplus confusus*, MS = *Melanoplus s. sanguinipes*, AP = *Arphia p. pseudonietana*, SB = *Spharagemon bolli*, SC = *Spharagemon collare*.

³Based on Gangwere (1962).

Table 2A. Size variation in defecating animals and their respective feculae.

Groups and Species Tested	Fecular N ⁴	Defecator N ⁴	Mean Fecular Length (mm)	Mean Defecator Length (mm)	Mean Variable Ratio ⁵
Males, all species	66	68	3.5121	19.5294	0.1806
Females, all species	90	90	4.4576	22.211	0.2252
Acrididae: Oedipodinae					
<i>Arphia p. pseudonietana</i>	26	28	4.7688	22.3214	0.2971
<i>A. p. pseudonietana</i> (males)	13	15	3.9938	21.6667	0.1845
<i>A. p. pseudonietana</i> (females)	13	13	5.5438	23.0769	0.4097
<i>Spharagemon b. bolli</i>	13	13	4.7015	25.7692	0.1828
<i>S. b. bolli</i> (males)	3	3	3.0867	21.0000	0.1463
<i>S. b. bolli</i> (females)	10	10	5.1860	27.2000	0.1937
<i>Spharagemon collare</i>	44	44	3.9768	21.0227	0.1892
<i>S. collare</i> (males)	27	27	3.6852	19.5556	0.1885
<i>S. collare</i> (females)	17	17	4.4400	23.3529	0.1902
Acrididae: Catantopinae					
<i>Melanoplus confusus</i>	48	48	3.7298	19.9375	0.1847
<i>M. confusus</i> (males)	13	13	2.6754	17.6154	0.1525
<i>M. confusus</i> (females)	35	35	4.1214	20.8000	0.1966
<i>Melanoplus s. sanguinipes</i>	25	25	3.7544	19.4000	0.1958
<i>M. s. sanguinipes</i> (males)	10	10	3.6340	18.3000	0.2007
<i>M. s. sanguinipes</i> (females)	15	15	3.8347	20.1333	0.1926

⁴N = number

⁵Variable ratio = defecator length/fecular length

Table 2B. Size variation in defecating animals and their feculae: by fecular type.

Types of Feculae ⁶	Fecular N ⁷	Defecator N ⁷	Mean Fecular Length (mm.)	Mean Defecator Length (mm.)	Mean Variable Ratio ⁸
1B	33	34	4.7521	21.0588	0.2918
1C	104	105	3.8388	21.0952	0.1815
BC	16	16	4.2306	21.2500	0.1995

⁶Based on Gangwere (1962).

⁷N = Number

⁸Variable ratio = defactor length/fecular length.

Table 3. One-way analysis of variance for variable ratios (feculae: animal) between various populations and groups.

Populations or Groups Tested	N 1	N 2	Mean Variable Ratio 1	Mean Variable Ratio 2	t Value	Probability ^a (P)	Significance
Type 1B: Type 1C	33	104	0.2918	0.1815	2.6303	0.001 ≤ P ≤ 0.01	(+)
Type 1B: Type BC	33	16	0.2918	0.1995	0.8692	0.4 ≤ P ≤ 0.5	(-)
Type 1C: Type BC	104	16	0.1815	0.1995	1.3585	0.1 ≤ P ≤ 0.2	(-)
Total ♂: Total ♀	66	90	0.1806	0.2252	1.3704	0.1 ≤ P ≤ 0.2	(-)
<i>Melanoplus confusus</i> ♂ & ♀:							
<i>Melanoplus sanguinipes</i> ♂ & ♀	48	13	0.1847	0.1828	0.1160	0.5 ≤ P	(-)
<i>Melanoplus confusus</i> ♂ & ♀:							
<i>Arphia p. pseudonietana</i> ♂ & ♀	48	26	0.1847	0.2971	1.6152	0.1 ≤ P ≤ 0.2	(-)
<i>Spharagemon b. bolli</i> ♂ & ♀:							
<i>Spharagemon collare</i> ♂ & ♀	13	44	0.1828	0.1892	0.4593	0.5 ≤ P	(-)
<i>Spharagemon b. bolli</i> ♂ & ♀:							
<i>Melanoplus sanguinipes</i> ♂ & ♀	13	25	0.1828	0.1958	0.8061	0.4 ≤ P ≤ 0.5	(-)
<i>Spharagemon b. bolli</i> ♂ & ♀:							
<i>Arphia p. pseudonietana</i> ♂ & ♀	13	26	0.1828	0.2971	0.8510	0.4 ≤ P ≤ 0.5	(-)
<i>Spharagemon collare</i> ♂ & ♀:							
<i>Melanoplus sanguinipes</i> ♂ & ♀	44	25	0.1892	0.1958	0.6266	0.5 ≤ P	(-)
<i>Spharagemon collare</i> ♂ & ♀:							
<i>Arphia p. pseudonietana</i> ♂ & ♀	44	26	0.1892	0.2971	1.4906	0.1 ≤ P ≤ 0.2	(-)
<i>Melanoplus sanguinipes</i> ♂ & ♀:							
<i>Arphia p. pseudonietana</i> ♂ & ♀	25	26	0.1958	0.2971	1.0511	0.2 ≤ P ≤ 0.3	(-)
<i>Melanoplus confusus</i> ♂:							
<i>Melanoplus confusus</i> ♀	13	35	0.1525	0.1966	2.7663	.001 ≤ P ≤ 0.01	(+)
<i>Melanoplus confusus</i> ♂:							
<i>Spharagemon b. bolli</i> ♂	13	3	0.1525	0.1463	0.4274	0.5 ≤ P	(-)
<i>Melanoplus confusus</i> ♂:							
<i>Spharagemon collare</i> ♂	13	27	0.1525	0.1885	3.2228	0.001 ≤ P ≤ 0.01	(+)
<i>Melanoplus confusus</i> ♂:							
<i>Melanoplus sanguinipes</i> ♂	13	10	0.1525	0.2007	3.2944	.001 ≤ P ≤ 0.01	(+)
<i>Melanoplus confusus</i> ♂:							
<i>Arphia p. pseudonietana</i> ♂	13	13	0.1525	0.1845	1.9570	0.05 ≤ P ≤ 0.10	(-)
<i>Melanoplus confusus</i> ♀:							
<i>Spharagemon b. bolli</i> ♀	35	10	0.1966	0.1937	0.1464	0.5 ≤ P	(-)
<i>Melanoplus confusus</i> ♀:							
<i>Spharagemon collare</i> ♀	35	17	0.1966	0.1902	0.4052	0.5 ≤ P	(-)
<i>Melanoplus confusus</i> ♀:							
<i>Melanoplus sanguinipes</i> ♀	35	15	0.1966	0.1926	0.2522	0.5 ≤ P	(-)
<i>Melanoplus confusus</i> ♀:							
<i>Arphia p. pseudonietana</i> ♀	35	13	0.1966	0.4097	1.8985	0.05 ≤ P ≤ 0.10	(-)
<i>Spharagemon b. bolli</i> ♂:							
<i>Spharagemon b. bolli</i> ♀	3	10	0.1463	0.1937	1.4102	0.1 ≤ P ≤ 0.2	(-)
<i>Spharagemon b. bolli</i> ♂:							
<i>Spharagemon collare</i> ♂	3	27	0.1463	0.1885	1.9201	0.05 ≤ P ≤ 0.1	(-)

Table 3 (Continued)

<i>Spharagemon b. bolli</i> ♂: <i>Melanoplus sanguinipes</i> ♂	3	10	0.1463	0.2007	1.9228	0.05 ≤ P ≤ 0.1	(-)
<i>Spharagemon b. bolli</i> ♂: <i>Arphia p. pseudonietana</i> ♂	3	13	0.1463	0.1845	1.1662	0.2 ≤ P ≤ 0.3	(-)
<i>Spharagemon b. bolli</i> ♀: <i>Spharagemon collare</i> ♀	10	17	0.1937	0.1902	0.1707	0.5 ≤ P	(-)
<i>Spharagemon b. bolli</i> ♀: <i>Melanoplus sanguinipes</i> ♀	10	15	0.1937	0.1926	0.05862	0.5 ≤ P	(-)
<i>Spharagemon b. bolli</i> ♀: <i>Arphia p. pseudonietana</i> ♀	10	13	0.1937	0.1845	0.3994	0.5 ≤ P	(-)
<i>Spharagemon collare</i> ♂: <i>Spharagemon collare</i> ♀	27	17	0.1885	0.1902	0.1336	0.4 ≤ P ≤ 0.5	(-)
<i>Spharagemon collare</i> ♂: <i>Melanoplus sanguinipes</i> ♂	27	10	0.1885	0.2007	0.8333	0.4 ≤ P ≤ 0.5	(-)
<i>Spharagemon collare</i> ♂: <i>Arphia p. pseudonietana</i> ♂	27	13	0.1885	0.1845	0.2746	0.5 ≤ P	(-)
<i>Spharagemon collare</i> ♀: <i>Melanoplus sanguinipes</i> ♀	17	15	0.1902	0.1926	0.1406	0.5 ≤ P	(-)
<i>Spharagemon collare</i> ♀: <i>Arphia p. pseudonietana</i> ♀	17	13	0.1902	0.4097	1.3532	0.1 ≤ P ≤ 0.2	(-)
<i>Melanoplus sanguinipes</i> ♂: <i>Melanoplus sanguinipes</i> ♀	10	15	0.2007	0.1926	0.4455	0.5 ≤ P	(-)
<i>Melanoplus sanguinipes</i> ♂: <i>Arphia p. pseudonietana</i> ♂	10	13	0.2007	0.1845	0.7541	0.4 ≤ P ≤ 0.5	(-)
<i>Melanoplus sanguinipes</i> ♀: <i>Arphia p. pseudonietana</i> ♀	15	13	0.1926	0.4097	1.2557	0.2 ≤ P ≤ 0.3	(-)

⁹p = the probability that a t value equal to that observed can be obtained by chance alone.

DISCUSSION

In terms of Gangwere's classification of feculae of orthopteroids (1962), the feculae of the species of Acrididae here investigated belong to types 1B (Fig. 1), 1C (Fig. 3), and BC (an intermediate between 1B and 1C) (Fig. 2). Type 1B feculae were described by Gangwere as typical of the Oedipodinae and Type 1C as typical of the Catantopinae. The former pellets, which are composed of somewhat aligned fibers, are comparatively regular and spindle-shaped in form, while the latter are always wrinkled, irregular, and unaligned. The occurrence of these three types is closely related to food. It was noted during the present study that grasses and their relatives consistently give rise to Type 1B feculae; Compositae and other "higher" forbs to Type 1C; and herbs phylogenetically intermediate between grasses and "higher" forbs to Type BC. These differences are interpreted as directly related to food composition.

Type 1B feculae are generally longer in comparison with length of defecator than are either types 1C or BC feculae. Type 1B feculae averaged approximately 29% of defecator length, Type 1C 18% and the intermediate BC 20% for the total number of animals measured (Table 2B, column 6).

The collective feculae of all females measured are larger than are those of the males, and this absolute difference in size is apparently related to the size differential between the sexes. Males of all species studied averaged 19.5294 mm. in body length, while the females




Fig. 1. Aligned fecula (Type 1B) of a grass-fed *Chortophaga viridifasciata* (Orthoptera: Acrididae: Oedipodinae).

Fig. 2. Poorly aligned fecula (Type BC) of a thistle-fed *Chortophaga viridifasciata*.

Fig. 3. Unaligned fecula (Type 1C) of a lettuce-fed *Chortophaga viridifasciata*.

averaged 22.211 mm. Thus, the feculae produced by the males are approximately 18% of the total body length, as opposed to those of the females, which approach 23% (Table 2A, column 6).

The larger species tend, in general, to have a greater average fecular length (Table 2A, column 6). However, the relative ratio of fecular length to length of body is relatively constant at 18 to 20% for all species, except *Arphia p. pseudonietana*, where the ratio is approximately 30%, possibly in response to that species' almost complete graminivory.

The males of a given species of acridid are always smaller than are the females, and with one exception have a correspondingly lesser ratio of fecular length to body length (Table 2A, column 6). However, this difference between the sexes is statistically insignificant (Table 3, column 8), ruling out the possibility of a physiological sexual dimorphism of the digestive system. *Melanoplus s. sanguinipes* proved to be the only exception to the rule of a higher ratio among the female sex. In this species, females average 1.8 mm. longer than do the males, but have a fecular length-body length ratio that is 1% lower than that of the males (Table 2A, column 6). This disparity, too, is statistically insignificant (Table 3, column 8).

The fecular length-body length ratio may now be considered as a function of food plant. Scrutiny of Table 2B (column 6) indicates that Type 1B feculae (composed of grasses and their allies) exhibit a higher ratio (29%) than do the forb-type 1C feculae (18%). Moreover, this disparity proves statistically significant (Table 3, column 8), which confirms the validity of the acridid fecal types proposed by Gangwere (1962).

One-way analysis of variance tests on the ratio of fecular length to animal length between various populations and groupings is also possible, and is here carried out. The results appear in Table 3. Based on that tabulation, males and females—regardless of species—show no significant differences in their ratios, and the individual species—regardless of sex tested—also show no significant differences. Indeed, of all possible combinations of species and sex tested, only those differences between the males of *Melanoplus confusus* and those of *M. s. sanguinipes*; the males of *M. confusus* and those of *Spharagemon collare*; and the males and the females of *M. confusus* prove significant (Table 3, column 8). These few significant differences could be related to some yet unstudied biological factor, but are more likely a result of the small sample size of *M. confusus* males.

SUMMARY AND CONCLUSIONS

The length of the acridid feculae here studied is in part related to size of defecator and in part to composition of food. The length of Type 1B feculae proves primarily a consequence of food source (being always formed of grasses), while that of Type 1C feculae is more closely related to size of defecator than to food. As expected, the intermediate Type BC feculae are intermediate in this respect. The conformation of acridid feculae (as opposed to pure length, above) is determined by the food composition. Thus, the tendency of acridids to select assortments of food species from only a few fundamentally different kinds of plants (*i.e.*, grasses, forbs, shrubs, etc.) leads inevitably to the production of a limited number of fecal types. These types were earlier named by Gangwere (1962), based on comparative length, alignment, and shape, and the validity of the acridid types that he proposed is statistically confirmed by the present study. It appears that, in general, the length and form of the feculae of grasshoppers is determined by the combined influence of defecator length and food plant composition, with food exerting the greater effect. It is most certainly *not* related to the taxonomic position of the defecator. Consequently, it is here concluded that the feculae of grasshoppers are unsuitable objects for use in systematics, except in those relatively few species of the world-wide fauna that are highly specialized in their food selection.

ACKNOWLEDGMENTS

The authors are indebted to several Wayne State University faculty and staff members who were of assistance in the preparation of this report. They include Dr. You C. Chen, Department of Pediatrics, School of Medicine, who assisted in the statistical analysis presented in the report; Professor Edwin G. Vann, Department of Biology, who reviewed the manuscript; Miss Marlene Nigosian, Department of Gynecology and Obstetrics, School of Medicine, and Miss Mary Talos, Department of Biology, who typed the several drafts of the manuscript; and Miss Rosemarie Rozsa, who drew the figures.

LITERATURE CITED

- Bhatia, D. R., and H. Chandra. 1967. Effect of food plants on colour, structure, and size of faecal pellets of the desert locust, *Schistocerca gregaria* (Forsk.). *Ind. J. Ent.* 29:102-103.
- Boldyrev, B. T. 1928. Biological studies on *Bradyporus multituberculatus*, F. W. *Eos* 4:13-56.
- Brown, A. W. A. 1937. Studies on the excreta of a grasshopper (*Melanoplus bivitattus* Say). *J. Exp. Biol.* 14:87-94.
- Frost, S. W. 1928. Insect scatology. *Ann. Entomol. Soc. Amer.* 21:36-46.
- Gangwere, S. K. 1960. The feeding and culturing of Orthoptera in the laboratory. *Entomol. News* 71:7-13, 37-45.

- Gangwere, S. K. 1962. A study of the feculae of Orthoptera, their specificity, and the role which the insects' mouthparts, alimentary canal and food-habits play in their formation. *Eos* 38:247-262.
- Gangwere, S. K. 1969. A combined short-cut technique to the study of food selection in Orthopteroidea. *Turtax News* 47:121-125.
- Gangwere, S. K., and E. Morales Agacino. 1964. The feculae ("feces") of some Orthoptera (*sens. lat.*) of Tunisia. *Entomol. News* 75:209-219, 242-251.
- Hodson, A. C., and M. A. Brooks. 1956. The frass of certain defoliators of forest trees in the north central United States and Canada. *Canad. Entomol.* 88:62-68.
- Morris, R. F. 1942. The use of frass in the identification of forest insect damage. *Canad. Entomol.* 74:164-167.
- Sardesai, J. B. 1969. Relationship of size of fecal pellets to larval molts in Lepidoptera. *Ann. Entomol. Soc. Amer.* 62:662.
- Scott, H. G. 1964. Pictorial key to the droppings of domestic rodents and cockroaches. U. S. Dept. Health, Education, and Welfare, Pub. Health Service. Washington.
- Simeone, J. B. 1964. The frass of northeastern U. S. powder posting beetles. *Proc. XIIth Int. Cong. Entomol., London (1964):707-708.*
- Weiss, H. B., and W. M. Boyd. 1950. Insect feculae. *J. N. Y. Entomol. Soc.* 58:154-168.
- Weiss, H. B., and W. M. Boyd. 1952. Insect feculae II. *J. N. Y. Entomol. Soc.* 60:25-30.

BRIEF NOTICES

ANNUAL REVIEW OF ENTOMOLOGY: VOLUME 15. Edited by Ray F. Smith and Thomas E. Mittler. Palo Alto, California: Annual Reviews, 1970. viii, 502 pp. \$10.00.

The 1970 volume includes the following papers: Recent advances in insect population dynamics; The ecology of stream insects; Insect mimicry; Evolution and taxonomic significance of reproduction in Blattaria; Interpretations of Quaternary insect fossils; The structure of arthropod chemoreceptors; Honey bee nutrition; Biosystematics of the Embioptera; Physiology of insect hearts; Biological rhythms in terrestrial arthropods; Recent advances in insect pathology; Mode of action of pyrethroids, nicotinoids, and rotenoids; Entomology of the cocoa farm; Pome fruit pests and their control; Ultralow volume applications of concentrated insecticides in medical and veterinary entomology; Mite transmission of plant viruses; Resistance of ticks to chemicals; and Mycoplasma and phytarboviruses as plant pathogens persistently transmitted by insects. There is an index of chapter titles in Vols. 6-15, arranged according to subject.

R. S. Wilkinson

ADVANCES IN INSECT PHYSIOLOGY: VOLUME 6. Edited by J. W. L. Beament, J. E. Treherne and V. B. Wigglesworth. London and New York: The Academic Press, 1969. x, 308 pp. \$13.00.

There are five contributions in the latest volume of this series: Frost resistance in insects; Neural control of firefly luminescence; Postembryonic development and regeneration of the insect nervous system; The biology of pteridines in insects; and Electrochemistry of insect muscle. Included are cumulative lists of authors and chapter titles for Vols. 1-6.

R. S. Wilkinson