

Author name: **Carly Hanson**

Faculty Sponsor: Jennifer Holt

Department: Chemistry

Title: *Spectroscopic Determination of Binding Constants of Modified β -cyclodextrins with Brooker's Merocyanine*

Abstract:

Host-guest complexation occurs when a guest molecule is held inside a host molecule through weak molecular forces. β -cyclodextrin, a host molecule, has a variety of uses and is often applied in pharmaceuticals. Binding constants of host-guest complexes of Brooker's merocyanine and various modified β -cyclodextrins (2-hydroxypropyl- β -cyclodextrin, sulfated β -cyclodextrin, and methyl- β -cyclodextrin) were studied using fluorescence and UV-Vis spectrometry to determine the strength of interaction between the host molecule and the guest molecule. By modifying the β -cyclodextrin, the effect of ionic forces, hydrogen bonding, and steric hindrance were compared. It was determined via fluorescence spectrometry that sulfated β -cyclodextrin had a binding constant of 38.3 M^{-1} , which was significantly lower than the determined binding constants 430 M^{-1} for β -cyclodextrin, 359.1 M^{-1} for 2-hydroxypropyl- β -cyclodextrin, and 194.6 M^{-1} for methyl- β -cyclodextrin. These results were confirmed via UV-Vis spectrometry, where the binding constants were 335.2 M^{-1} for 2-hydroxypropyl- β -cyclodextrin

and 178.4 M^{-1} for methyl- β -cyclodextrin. These results were in agreement with fluorescence data. The much lower binding constant of sulfated β -cyclodextrin could be caused by a combination of ionic forces and steric hindrance. A comparison of these results to the theoretical models will lead to verification of the effect of these forces on binding.

About the author:

Carly Hanson is a junior Chemistry major and German minor.