

2022

Inflammatory bowel disease: pathogenesis, diagnosis and current therapeutic approach

Adrian Silaghi

St. Pantelimon Clinical Emergency Hospital, Department of General Surgery, Bucharest, Romania

Vlad Denis Constantin

Carol Davila University of Medicine and Pharmacy, Department of General Surgery, Bucharest, Romania, constantindenis@yahoo.com

Bogdan Socea

Carol Davila University of Medicine and Pharmacy, Department of General Surgery, Bucharest, Romania

Petrișor Banu

Carol Davila University of Medicine and Pharmacy, Department of General Surgery, Bucharest, Romania

Vladimir Sandu

See next page for additional authors

Follow this and additional works at: <https://scholar.valpo.edu/jmms>



Part of the [Gastroenterology Commons](#), and the [Surgery Commons](#)

Recommended Citation

Silaghi, Adrian; Constantin, Vlad Denis; Socea, Bogdan; Banu, Petrișor; Sandu, Vladimir; Andronache, Liliana Florina; Dumitriu, Anca Silvia; and Paunica, Stana (2022) "Inflammatory bowel disease: pathogenesis, diagnosis and current therapeutic approach," *Journal of Mind and Medical Sciences*: Vol. 9: Iss. 1, Article 6.

DOI: 10.22543/7674.91.P5677

Available at: <https://scholar.valpo.edu/jmms/vol9/iss1/6>

This Review Article is brought to you for free and open access by ValpoScholar. It has been accepted for inclusion in *Journal of Mind and Medical Sciences* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

Inflammatory bowel disease: pathogenesis, diagnosis and current therapeutic approach

Authors

Adrian Silaghi, Vlad Denis Constantin, Bogdan Socea, Petrișor Banu, Vladimir Sandu, Liliana Florina Andronache, Anca Silvia Dumitriu, and Stana Paunica

Inflammatory bowel disease: pathogenesis, diagnosis and current therapeutic approach

Adrian Silaghi¹, Vlad Denis Constantin^{1,2*}, Bogdan Socea^{1,2}, Petrișor Banu^{1,2}, Vladimir Sandu¹, Liliana Florina Andronache³, Anca Silvia Dumitriu⁴, Stana Paunica⁴

¹ST. PANTELIMON CLINICAL EMERGENCY HOSPITAL, DEPARTMENT OF GENERAL SURGERY, BUCHAREST, ROMANIA

²CAROL DAVILA UNIVERSITY OF MEDICINE AND PHARMACY, DEPARTMENT OF GENERAL SURGERY, BUCHAREST, ROMANIA

³CAROL DAVILA UNIVERSITY OF MEDICINE AND PHARMACY, BUCHAREST, ROMANIA

⁴CAROL DAVILA UNIVERSITY OF MEDICINE AND PHARMACY, DAN THEODORESCU HOSPITAL, BUCHAREST, ROMANIA

ABSTRACT



Inflammatory bowel disease is a group of conditions of unknown etiology, represented by Crohn's disease and ulcerative colitis and characterized by the presence of intestinal wall inflammation. From the first cases described to date, several studies have been performed to elucidate the cause of inflammatory bowel disease. Generally, the genetic factors predispose to their occurrence while epigenetic and environmental factors trigger them. Genetic factors are mutations of the genes involved in the response and recognition of immune cells to different pathogens. The most studied epigenetic and environmental factors are smoking (which predisposes to Crohn's disease and can be protective for ulcerative colitis), lack of vitamin D, a diet rich in sugars and low in flavonoids and fibers. The diagnosis is usually established by endoscopy and biopsy. Recent technologies can perform live biopsies such as endocytoscopy or confocal laser endomicroscopy, with an accuracy of 100% compared to classical methods. Therapy involves several classes of drugs, preferably in association with diet and lifestyle changes. In case of complications or in non-responsive diseases, surgery must be considered, as documented for ulcerative colitis that can be cured by removing the entire colon. The purpose of this review is to present recent findings on pathology, as well as modern diagnosis and treatment methods for IBD.

Category: Review

Received: November 14, 2021

Accepted: January 21, 2022

Published: April 10, 2022

Keywords:

inflammatory bowel diseases, pathogenesis, diagnosis, therapeutic approach

*Corresponding author:

Vlad Denis Constantin,

Carol Davila University of Medicine and Pharmacy, St. Pantelimon Hospital, Department of General Surgery, Bucharest, Romania

E-mail: constantindenis@yahoo.com

Introduction

Inflammatory bowel disease (IBD) is a group of diseases characterized by the inflammation of the intestinal wall. This can include a layer (mucosa) or all intestinal layers, with severe complications such as perforations, stenosis, bleeding, all of which having a major impact on the patients' life quality. The economic impact is major due to the chronic treatment, high hospitalization rates and recurrence [1-3]. This group of diseases includes Crohn's disease/ CD, characterized by the presence of transmural inflammation in any segment of the digestive tract (clinically expressed as diarrhea with steatorrhea, weight loss, fever, abdominal pain, etc.), and ulcerative colitis/ UC, pathology located mainly in the colon - most often rectally, where only the mucosal layer is affected (the main symptom being the occurrence of bloody stools) [4-7]. A subtype of inflammatory bowel disease has also been described, which borrows the characteristics from the CD and UC, or which cannot fall into either of the two major

categories. It is classified as unqualified inflammatory disease [8,9]. Their incidence has increased in recent years, occurring in both developed and developing countries, with a different distribution by sex. Crohn's disease is more common in women and is associated with other autoimmune diseases/manifestations (irises, irido-cyclitis, autoimmune thyroiditis, etc.), as well as more frequent digestive complications (perforation with abscess, stenosis and fistula formation) [10].

The etiology of the disease remains unknown today, most likely due to the interaction between immunological genetic factors (imbalance between anti-inflammatory and pro-inflammatory factors, with a predominance of the latter). Other factors are represented by microbiotics (patients with IBD have a microbial flora with a prosthesis activity 9-10 times higher than that of healthy individuals), smoking, a low-fiber diet and a rich processed food diet/refined food [11,12]. The diagnosis of patients with inflammatory bowel disease is established through colonoscopy (which can reveal the appearance of the

mucosa, the presence of ulcerations, stenosis and, at the same time, it can procure bioptic material for histopathological examination), computed tomography or MRI scanning [13-15]. The current diagnosis by serologic markers is possible by fecal calprotectin that can distinguish between irritable bowel syndrome and inflammatory bowel disease, with the mention that the fecal level can be influenced by several factors (sex, age, fiber, the presence of blood in the stool, etc.) [16,17].

The treatment of inflammatory bowel disease has undergone significant changes since the discovery of the pathological entity so far, with several classes of drugs being available for administration (amino salicylates, corticosteroids, immunological and molecular treatment). In case of complications, surgical intervention is also required. A breakthrough was the discovery of the treatment that can interfere with the TNF pathway, drugs that can induce the long-term remission. The new therapeutic approach is based on the direct influence of the cellular mechanisms that lead to the gastro-intestinal lesions, the final goal being the healing of the mucosa (elimination of inflammation with the appearance of normal mucosal structure) [18-21].

The purpose of this review is to provide an update on the literature on the causes of IBD, as well as modern methods of diagnosis and treatment.

Discussion

Pathogenesis

The occurrence of IBD is determined by the interaction between several factors, such as: genetic, immunological, environmental and microbiota factors.

Genetic factors

To date, 163 genes have been discovered in patients with inflammatory bowel disease, of which 30 genes are common for UC and CD, suggesting the existence of a common pathological pathway [1]. The NOD-2 gene encodes a protein that functions as an intracellular receptor and it structurally resembles the peptidoglycan on the surface gram-positive and gram-negative bacteria. It helps the immunogenic cells in the process of autophagy, which can control the bacterial replication and modulate the innate and adaptive immune system. It has been observed that mutations of NOD-2 gene can predispose to the occurrence of CD [2]. In the study of the pathophysiology of IBD, it was discovered that autophagy plays a central role, because it can control the intracellular homeostasis by degrading cytosolic elements and organs [3], ATG16L1 also appears to be involved in the intracellular process, and T300A point mutation may predispose to CD. IRGM is a protein which belongs to the p47 immunity-related GTPase family, its expression on the epithelial and dendritic cells being often decreased in CD patients [3,4].

Genetic sequencing methods have been used to identify the IL-23 receptor, in order to establish a possible association between its mutations and CD [5]. IL-23 is a pro-inflammatory cytokine that is involved in the generation of T-helper-17 (Th17) cell, being related to the pathogenesis of IBD (with gene loci: IL-23R, IL-12B, JAK-2 and STAT-3) [6,7]. In addition, genetic defects of IL-10, CARD-9, IL-1R2 REL SMAD3 and other immune markers also predispose to IBD [4].

Environmental factors

Environmental factors are involved in the onset and evolution of IBD, among the most studied being: smoking, a diet rich in processed foods, a diet low in fiber and flavonoids, drug use, and psychological factors [8]. Smoking is the most investigated environmental factor, smokers having less often and less severe episodes of UC [9,10], while chronic disease is a risk factor for postoperative IBD [11].

The chronic administration of non-steroidal anti-inflammatory compounds/ NSAIDs and the large-scale use of antibiotics have long been considered risk factors for the development of IBD. It has been shown that the use of aspirin, regardless of the duration of administration, rhythm and doses, does not trigger nor impair the condition of patients with IBD, while the long duration of the treatment and high doses of NSAIDs have been associated with relapses [12]. Chronic antibiotics administration can cause changes in the bacterial flora, predisposing to infections with potentially pathogenic bacteria which further lead to colitis. The dose and the period of administration that would lead to the occurrence of IBD are not yet determined, remaining to be established in future studies [13].

The incidence of IBD has increased proportionally with the level of economic and industrial development, the confirmation being based on epidemiological and ecological studies. It is suggested that air pollution can lead to the increased incidence of IBD (the pathophysiological mechanism being the increase in the plasma levels of cytokines and circulating cells) [14-16], while the increased level of SO₂ and NO₂ secondary to the burning of fossil fuels can lead to frequent and severe relapses [17].

Vitamin D plays a role in the pathogenesis of IBD, its deficiency being considered a risk factor for their occurrence [18]. The World Health Organization (WHO) classifies the low intake of vitamin D into deficiency (the plasma level of 25-hydroxyvitamin D [25(OH)D] being below 10ng/ml) and insufficiency (in which the level varies between 10-25ng/ml) [19]. In the digestive tract, Vitamin D maintains the integrity of the enteric mucosa, by increasing the number of tight and adherent junctions [20], E-cadherin and also some components of tight junctions (such as occludin and claudin) [21]. Another effect of Vitamin D is to decrease the kinase activity which can

degrade the myosin chain (due to its binding to the nuclear receptor) and prevents the degradation of the tight junctions [22]. Studies concerning the number and the activity of vitamin D receptors in patients with IBD are controversial. Thus, there is only one study that finds a lower number of receptors for vitamin D in subjects with active IBD than in those who did not suffer from it [23], while other two studies do not find statistically significant data between the groups, although they find different numbers of receptors between the normal and the affected mucosa [24,25]. In patients with active IBD, there is a decrease in the expression of Vitamin D receptors, secondary to the mucosal inflammation mediated by TNF α and over-expression of CYP27B1 [26]. Binding the vitamin to the receptor can reduce the activity of claudin-2 (the protein involved in non-specific trans-cellular permeability and apoptosis), thus being a protective factor for IBD [27]. This was confirmed by Garg et al., who administered 10,000 IU of vitamin D daily; after 12 weeks, the serum levels of this vitamin in patients varied between 40-50ng/ml, being accompanied by a reduction of the disease activity on the clinical and endoscopic scale [28].

The consumption of flavonoids in food is considered a protective factor against inflammatory bowel disease, as they have antioxidant properties by decreasing the amount of free radicals (causing changes in the microbiota), thus protecting the mucosal barrier [29]. The antioxidant properties of flavonoids are favorable in inflammatory bowel disease, where the free radicals of oxygen are overproduced secondarily to myeloperoxidase (MPO) and NADPH system activity [30]. In vivo studies have shown that most flavonoids reduce colonic MPO levels by lowering free radical levels, which can be considered a marker for leukocyte invasion and inflammation [31]. Daidzein, hesperidin and naringenin flavonoids are found in citrus fruits, bitter fruits and tomatoes, having a protective capacity on the enteric mucosa by maintaining the integrity of tight junctions. It is currently considered that the hyper permeability of the mucosa is the initiating factor of inflammatory bowel disease [32]. Flavonoids are also involved in modulating the immune system at the intestinal level, decreasing the amount of pro-inflammatory cytokines including TNF α , IL-6, IL-1b via Epigallocatechin gallate [33].

Quercetin is a flavonoid found ubiquitously in fruits and vegetables, and causes a decrease in the amount of macrophages and neutrophils infiltrated in the colon mucosa of patients with CD or UC, thus decreasing the activity of the disease [34]. In patients with IBD, severe dysbiosis occurs with a decrease in the biodiversity of the colon, the main species found being Proteobacteria [35]. This aspect has been identified in mice with induced colitis; addition of Quercetin as a supplement appears to be a favorable factor, by enhancing the biodiversity in the colon and reducing the inflammation [36].

Immunological factors

In the pathogenesis of inflammatory bowel disease, there are changes in the innate and acquired immune system that led to chronic inflammation, secondary to the impaired barrier function of the mucosal layer, changes in the intestinal microbiota and disorders of autophagy [4]. Innate immunity is the first line of defense, being nonspecific and causing a rapid immune response. Its main role is to recognize various antigens through the NOD and TLR receptors, which are found on the surface and in the cytoplasm of various immune cells (neutrophils, dendritic cells, monocytes, etc.) [37]. In patients with IBD, there is an alteration of these two types of receptors. The mutation NOD2 3020insC 2 predisposes to CD. The mechanisms presumed to be involved are the lack of response of the receptor to bacterial lipopolysaccharides [38]. In addition, the lack of activation of the NF-Kb pathway can reduce the antibacterial response and will lead to the appearance of bacterial overpopulation [39,40]. Yet, the lack of inhibition of the RLR2 pathway leads to an excessively immune response modulated by T helper cell 1(Th1) [41].

Interleukin-23 plays an important role in the function of both the innate and adaptive immune system, intervening early in the response to a pathogen; its receptor polymorphism is often associated with the presence of both UC and CD [4].

The acquired immune response is highly specific and it requires a long time to perform the function. Through the Th1 cell, large amounts of IL-12 and IFN- γ are produced, and through Th2 helper cells, IL-4, IL-5 and IL-13 are released [42]. In CD, there are high levels of IL-2 and IFN- γ secondary to increased Th1 cell activity; in contrast, UC has high serum concentrations of IL-13 secondary to natural killer-T cell hyper activation [43,44]. The classical assumption in CD is that the immune response is mediated by Th1 cells, while in UC by Th2 cells [45]. In vivo studies have shown that there are similar levels of INF in both pathologies [46] and that low levels of IL-13 (with anti-inflammatory function) are similar [47]. Bernardo et al. identified a mixed cytokine profile with predominancy of IL-6 and the absence of IL-13 in the mucous cell culture of patients with IBD, which therefore leads to a change in the classical conception [48]. Th17 cells are capable of synthesizing large amounts of IL-17, IL-21 and IL-22, which are the result of the concomitant action of IL-6 and transforming growth factor β . IL-17A can be found in large amounts in both CD and UC [49]. The definite role of these cells in the pathogenesis of IBD remains to be investigated, especially in that they express surface receptors for IL-23.

Clinical aspects of IBD

Intestinal manifestations of IBD

The characteristic of **Crohn's** disease is the ubiquitous involvement of the digestive tract from the oral cavity to the rectum and anal canal, which can take great variability.

In classical semiology, a relatively specific pattern is described - the appearance of abdominal pain with colic, weight loss and watery diarrhea [50]. Frequently, abdominal pain occurs prior to the diagnosis of the disease for several years, its location being in the lower-right quadrant because the terminal ileum is affected. The character of the pain is colicky and the intensity is medium to severe. It can often be confused with appendicitis [51]. Diarrhea may be watery, sometimes with the presence of bloody streaks, more frequently in case of rectal damage [52]. Bleeding can also occur many years before the diagnosis of the disease, with 1-2% of cases with a vital threat due to high flow [53]. Weight loss is the third major symptom with a multifactorial cause, presence of chronic inflammation leading to malabsorption and anorexia.

Chronic disease is often associated with the presence of fistula and stenosis. The most frequent type of fistula is between the separate segments of the jejunum, defined as entero-enteral fistula. Most of them are asymptomatic; if the diameter of fistula enlarges, the process can lead to an accelerated intestinal transit with the occurrence of watery diarrhea. The leakage of intestinal fluid through the vagina shows the presence of an entero-vaginal fistula. In a few cases, there may be abnormal communication between the digestive tract and the urinary tract or even the skin. The processes located in the digestive tract can be manifested by colic pain, abdominal bloating or early satiety. For those placed in the stomach or the duodenum, symptoms vary depending on the location [51]. In one third of the cases, patients with CD have anal suffering through the presence of fistulas, abscesses and fissures [54]. The presence of anal fistulas is suggestive for a special clinical form, as these patients have relapses with high and severe recurrence rates that require prompt therapeutic intervention [55].

Regarding **ulcerative colitis**, the symptoms are more stable, the spread of the disease being exclusively in the colon. The time until the diagnosis being much shorter than in the case of CD. The main clinical manifestation of UC is marked by diarrhea (that occurs most often postprandial and at night), its severity correlating with the severity of the inflammation. Regarding the extent of the disease in the colic region, if the proximal part of the colon is affected, a large quantity of mucus and blood will come along with the stool [51]. Massive bleeding with hemodynamic instability is rare, and can be found in cases of ischemic colitis or toxic mega colon, accounting for 15% of the patients [56].

Other symptoms present in patients with UC are abdominal colic pain along with other general symptoms such as asthenia, fatigue and fever. With the long-term evolution of the disease, the appearance of strictures and stenosis can be seen, which can lead to colonic constipation. It can occur in the distal locations of the disease, when the stool is associated with lower digestive bleeding [57].

Extra-intestinal manifestations of IBD

IBD is actually a veritable systemic disease, distant manifestations occurring in addition to the local lesions. Their incidence varies between 6.2-46.6%, thus affecting the quality of life of the patient [58]. It is considered that if one extra-intestinal manifestation appears, the patient is likely to develop other systemic manifestations over time. The way they appear in relation to the time of diagnosis of inflammatory bowel disease can be variable. That is, before the onset of digestive symptoms (8.7-10% of patients, according to the study), concomitantly or in the first month after diagnosis (about 25% of the cases) and in the next two months after the diagnosis for the rest of the patients [59].

The course of extra-intestinal manifestations is not well documented. It is considered that antigenic structures presenting similarities with epitopes of cells in the skin, synovial capsule or periarticular tissue, can pass through the intestinal mucosa triggering a local immune response [60-62]. Patients with CD presenting HLA-A2, DR-1, DQw5 have a significant predisposition to develop extra-intestinal manifestations, which are similar to patients with UC presenting HLADR103 [63]. In the major histocompatibility complex, there are certain subtypes that predispose to several extra intestinal manifestations. For example, HLA-DRB1*0103, HLA-B*27 and HLA-B*58 predispose to articular, dermal and ocular diseases [64,65].

Musculoskeletal manifestations are the most common form of injury in IBD patients (small and axial joints are often affected), being present in approximately 40% of the cases. These include arthritis and peripheral arthritis where there is no joint damage, as opposed to psoriatic arthritis or rheumatoid arthritis [66,67]. This form of arthritis is divided into two subtypes. Type 1 of polyarticular condition usually is pauciarticular, with fewer than 5 joints being affected and associated with HLA-B27, B35, and HLA-DR103 [68]. Large joints (especially the knee) are damaged, and the treatment administered for intestinal manifestations can improve and relieve the joint pain [67]. Type 2 polyarticular condition is characterized by the affliction of more than 5 joints, which does not necessarily correlate with the evolution of the digestive disease, but is instead associated with uveitis [69]. In general, arthritis responds to the NSAIDs (especially COX₂), with the metacarpophalangeal joint being most commonly involved [67,70].

Axial atrophies include ankylosing spondylitis and sacroiliitis, which are less common in incidence (only 25% of patients presenting such symptoms) [71]. In general, men are more likely to develop such arthropathies, especially HLA-B27 positive patients [72]. The evolution is generally progressive and causes significant skeletal damage without treatment [67]. Sacroiliitis usually occurs in HLAB27 negative patients and often causes chronic pain and incapacity of work, which lead to decreased work

capacity and quality of life [73]. In the case of axial arthropathy, the recommended treatment is methotrexate in combination with sulfasalazine and azathioprine, or anti-TNF antibodies of different generations in refractory cases [74-76].

Mucocutaneous damage occurs in about 15% of the patients, the main forms of manifestation being erythema nodosum, pyoderma gangrenosum, sweet syndrome and lesions of the oral mucosa [77].

Erythema nodosum is a nodular skin lesion (with dimensions varying between 0.5-4.5 cm) that is found on the anterior surface of the lower limbs and less often on the trunk or face; it usually appears in patients with CD. Healing is most often possible without scarring. The diagnosis is made clinically based on the characteristic appearance [78]. Regarding the treatment, they respond well to the systemic and/or the local treatment with corticosteroids (CSs) [79]. In severe cases that do not respond to the treatment, the suspicion of another cause of nodules (such as bacterial infections with pseudo tuberculosis, enterocolic, sarcoidosis, tertiary syphilis or drug reactions) may be considered. After their exclusion, there is an upgrade of the treatment with the administration of CSs in immunosuppressant doses, or the treatment with anti TNF. It is worth mentioning that the effect of Infliximab and Adalimumab is superior to other anti-TNF drugs [80-82].

Pyoderma gangrenosum occurs by the presence of a nodule or morbillus that grows rapidly in size and affects the adjacent skin tissue, with the development of a purplish ulcer at the edges with the elimination of sterile pus [83]. The size of the lesions varies from a few centimeters to the surface of a limb. Elective locations are on the extension surface, near the postoperative stoma, but they can occur anywhere on the body surface [84]. It is often associated with UC, with damage of the entire colon and affecting especially woman [85]. The declassifying factor is represented by skin trauma, including venous punctures or biopsies [86].

Regarding the treatment of pyoderma gangrenosum, along with the remission of the colonic manifestations, its remission also appears. The mild forms respond to the local injectable treatment with CSs, the local wound cleaning with wet pads [87]. Sulfasalazine, dapsone or immunomodulatory drugs are administered in refractory cases, in the form of azathioprine, cyclophosphamide, methotrexate, tacrolimus and anti-TNF agents. Most of the time, healing occurs with residual scars [87-89].

Sweet's syndrome is a rare manifestation associated with IBD, characterized by a papular-scaly rash on the hands, feet, torso, or face being associated with fever, arthropathy and ocular symptoms [90,91]. At the histopathological examination, large infiltrates with neutrophils appear. It is often associated with chronic diseases and possible

malignancies [67]. The treatment consists in the topical or systemic administration of CSs [92].

The lesions at the level of the oral cavity appear in approximately 10% of the IBD cases, determining the appearance of periodontitis, foot-and-mouth disease, or in severe cases pyosomatitis vegetans (characterized by pustular rashes anywhere on the surface of the oral mucosa) [93]. The treatment requires the administration of local CSs and/ or an antiseptic treatment [94].

Ocular manifestations occur in 2-5% of the patients [68,95], being more common in patients with CD than in UC, especially in subjects younger than 40 years [58,68]. Generally, uveitis and episcleritis (which affects the anterior chamber of the eye) can advance to the retina and the choroid, this process being known as panuveitis. Without a prompt diagnosis, uveitis can lead to vision loss. The treatment with topical steroids can be life-saving; in cases of refractory disease, cyclosporine A can be used [67]. The treatment with Infliximab appears to be the best solution for the patients with uveitis, sacroiliitis and CD [96].

Hepato-biliary damage in IBD consists of stricture and dilatation of the bile ducts, which can lead to intrahepatic lithiasis, cholecystitis, granulomatous hepatitis or hepatic steatosis [97]. It occurs more often in patients with UC than in CD [98]. The dilatations and strictures in the bile ducts appear before the diagnosis of CD and, once formed, they lead to altered bile leakage with the appearance of cholestasis and biliary cirrhosis [99]. The treatment of the underlying disease does not improve the dilatation and the stenosis. However, the administration of ursodeoxycholic acid improves liver functioning. In the case of extra hepatic strictures, the insertion of stents by ERCP procedure can be performed. In the final stage of the disease, a liver transplant is necessary [100].

The diagnosis of IBD

The diagnosis of inflammatory bowel disease is made by correlating the clinical signs and the symptoms with the paraclinical investigations (abdominal ultrasound, lower digestive endoscopy, CT, MRI, etc.) and serological tests, which are used both as a method of initial diagnosis and to monitor the disease evolution.

The endoscopic procedures

Endoscopic procedures play a major role in the diagnosis of inflammatory bowel disease. They are used for diagnosis/ biopsies (to differentiate between CD and UC) on the one hand, and for monitoring the disease evolution (response to treatment and complications) on the other hand [101]. In patients with IBD upon the initial evaluation, colonoscopy is performed in order to evaluate the terminal ileum with the collection of 2 biopsies from 5 different places, including the rectum and the terminal ileum [102]. It is a simple procedure with minimal side

effects, which must be limited on patients with severe forms of colitis or with toxic megacolon. In such cases, it is possible to perform only rectosigmoidoscopy to obtain tissue for the histopathological examination [103]. The classical aspect of UC is the loss of vascularity, erythema of the mucosa with granular and friable appearance, ulcers, erosions and pseudo-polyps. This aspect evolves from the rectum to the proximal level where the intensity gradually disappears [104]. In CD, the classical lesions are represented by the presence of discontinuous ulcers, which give the appearance of paving stone/ cobblestone [105].

In the last decade, the endoscopic equipment has been developed to highlight the presence of inflammation in the colonic and intestinal mucosa, being used to better diagnose/ evaluate the activity of the disease compared to classical endoscopy [106]. Dye-chromoendoscopy uses staining agents (methylene blue/ indigo carmine) that are applied to the mucosa, thus increasing the detective rate of any changes [107]. This investigation helps detect the early and various adenomas that could become malignant, secondary to the inflammation of the mucosa. The sensitivity of the endoscopic examination is correlated with the histopathological data [108,109].

Dye less-chromoendoscopy uses several working systems such as narrow band imaging (a narrow spectrum of light that is absorbed by hemoglobin from the vessels), thus leading to a better highlighting of the vascularization at the level of the organ of interest [110,111]. Compared to classical endoscopy, it is much better in assessing the vascularity which appears to correlate with histological inflammation [112]. The shape of the blood vessels can predict the risk of recurrence. Patients with vessels like bare branches have active disease, while in subjects presenting a honeycomb pattern the disease is often inactive [113]. Other dye-less chromoendoscopy technologies are represented by I-scan, blue light imaging, linked color imaging [110]. The correlation between the histopathological evolution and the endoscopic appearance on the I-scan has been recently investigated. The results are conclusive, with the OE score being largely correlated with two histological activity scores (Robarts histopathology index and ECAP for UC) [114]; similar results have been found in CD [115].

Endocytoscopy is a diagnostic method that allows the in vivo evaluation of the appearance of the mucosa; by obtaining virtual biopsies, it can reach up to 50µm in size [106]. It also correlates 100% with the activity investigated by classical methods (on Riley score) [116]. Recent studies have concluded that endoscopy is able to differentiate between patients with active disease and subjects in remission, without taking biopsies [117]. Confocal laser endomicroscopy is an endoscopic evaluation method that allows the evaluation tissues up to 250 µm [118]. Patients with remitted UC may show distorted crypts, inflammation

and abnormal vascular pattern [119]. In patients treated with TNF-alpha inhibitors, the number of affected crypts, the degree of vascular permeability and the degree of inflammation compared to the Gupta score could be assessed, with similarities in 95% of the cases [120].

Upper digestive endoscopy is useful in the evaluation of patients with CD, which can affect the upper portion of the digestive tract in 16% of the cases [121]. In such cases, two biopsies should be obtained from the esophagus, stomach and duodenum when the damage is suspected [122]. The results are characterized by the presence of ulcers, strictures, fistulas and erythema. These damages may be simultaneous or may occur later, when the manifestations appear in the distal part of the digestive tract [123]. In the pediatric population, it is recommended that the initial assessment be made regardless of the onset of the disease and whether upper digestive endoscopy is performed or not [124].

Small bowel capsule endoscopy is a sensitive method of highlighting small bowel damages, being recommended for patients with CD [125]. Since its introduction, technological leaps have been made in terms of image resolution and capsule size, which has led to their widespread use [126]. It is a useful investigation especially in the follow-up of patients, because it is able to diagnose patients with intestinal ulcers, thus allowing a step forward in terms of treatment and preventing the risk of intestinal stenosis and fistulas. At the same time, it evaluates the IBD evolution and response to treatment, allowing thus an anticipation of 6 months before the appearance of a new active form of the disease [127].

Artificial intelligence refers to the ability of computers to function similarly to the human mind, which can be useful for the endoscopic diagnosis of IBD [128]. A Google-based computer aided diagnosis (CAD) system was used to identify the normal mucosa and the presence of mucosal healing (Mayo score 1 for 114 patients with UC), this system recognizing more than 90% of the processed images [129]. Furthermore, with the help of artificial intelligence and based on the appearance of red areas on the endoscopic examination (given by the vascular pattern), the same team developed a red density score. This score correlates with endoscopic (Mayo) and histological activity data, which suggest that the system may intervene in the subjectivity of the investigator in the evaluation of the disease [130].

Imaging evaluation

In the IBD, a large number of imaging methods are used, such as: abdominal plain radiography, barium contrast studies, computed tomography, ultrasound and MRI.

Plain radiography is a simple, easy and inexpensive investigation, being used especially in emergency departments for the occurrence of IBD complications. The

digestive perforation induces pneumoperitoneum, while stenosis on intestinal ulcers appears with hydro-aerial levels and marked distension of the loops above the stenotic area [131]. This is most often completed by a spiral assessment of the abdominal floor such as CT/ MRI or abdominal ultrasound [132]. Fluoroscopic imaging is a method of assessing the digestive tract damages in CD and can be used to investigate both the small intestine and the colon. Regarding the upper digestive tract, the method of investigation has recently been replaced by CT, MRI or enterocapsule [133]. In order to highlight the lesions, it is necessary to mount a nasal-jejunal probe and to insufflate the air or methylcellulose (with the formation of a characteristic double contrast), the appearance of thickening of the intestine suggesting the presence of stenosis [134]. Contrast-based enemas can also be performed, highlighting erosions, ulcers or irregularities in the colon, with the mention that suggestive images result only in the presence of double contrast [135].

Computed tomography

Computed tomography (CT) is one of the first methods of evaluating patients with IBD, due to its availability and effectiveness. It can diagnose both the local complications of diseases (fistulas, abscesses, etc.) and the extraintestinal manifestations (pancreatitis, cholangitis, sacroiliitis, etc.) [131]. By administering intravenous contrast, several changes can be observed, such as the thickening of the mucosa and the mucous layer, the presence of adenopathy/ large nodules obvious in the middle of the affected region or Comb-sign, which is characterized by a turgescence vasa recta [136]. The CT can also be used to evaluate the activity of CD, with a sensitivity of 81% and a specificity of 88%, being able to detect strictures or damage to the peri-intestinal tissues [137]. Regarding UC, the results are moderate, the tomographic evaluation being less correlated with the activity of the disease (a sensitivity of 74%) [138]. It is more useful in detecting acute complications of the disease such as toxic megacolon, digestive perforations, etc. [139]. However, CT has limited utility in monitoring disease activity in young patients with IBD, as the effect of ionizing radiation may increase the risk of colon cancer by 7% (in young people) and 16% (in the elderly) [140].

Magnetic resonance imaging

MRI is an imaging diagnostic method that uses non-ionizing radiation. Its availability has recently increased, but it is not frequently used due to the extended time required for a complete evaluation [131]. It can be used to evaluate the small intestine and the perineal area (highlighting lymphadenopathy, fistulas and fissures with different trajectories), thus becoming the gold standard in identification of periintestinal complications [141]. Other signs observed by imaging methods may reveal the parietal thickening of the small intestine, hyper vascularization, abscesses and stenosis [142].

In IBD, the evolution can be investigated by MRI, with a sensitivity and specificity of over 80% in CD [143], and over 85% in UC [144]. An activity score was developed by Rimola-Magnetic Resonance index of Activity/ MaRIA score, in which several parameters are included (intestinal wall thickness, contrast loading, edema and ulceration) [145]. The MaRIA score is obtained by quantifying these parameters at the ileal level, colon and rectum. These results are correlated with Crohn's Disease Endoscopic Index of Severity (CDEIS), so the MaRIA score correlates with the disease activity and evolution (a score > 7 appears in the active medium forms, while a score > 11 in the severe forms of the disease) [145,146].

The treatment of IBD

Several medications are currently available for therapy of IBD, such as: amino salicylates, immunomodulators, corticosteroids, biological treatments and small molecules.

Aminosalicylates

Aminosalicylates are represented by sulfasalazine and other 5-aminosalicylic acid derivatives. Sulfasalazine is a compound that has been used for over 80 years in the treatment of IBD [147]. Its mechanism of action is related to the arachidonic acid system, which causes a decrease in the amount of leukotrienes and prostaglandins, the inactivation of reactive and nitrogen reactive species, and the activation of the aryl hydrocarbon receptor pathway (which determines the activation of the tissue growth factor with the accentuation of the tissue recovery) [148,149]. The action of aminosalicylates in UC is documented, as it can induce the remission of the disease regardless of the rate of administration (1 or 2 doses daily) [150]. At the same time, it represents a protective factor for malignant complications, reducing the risk of colon cancer by 75% [151]. Regarding CD, studies have shown that the oral administration of aminosalicylates does not induce remission nor it maintains it [152]. However, their administration for more than 1 year has led to decreased medical costs of hospitalization for severe forms or surgery [153]. Mesalamine, when administered in high doses, induces remission of the disease (in patients with moderate and severe forms of CD) and prevents the administration of steroids [154].

Adverse reactions caused by aminosalicylates are moderate (headaches, abdominal pain and diarrhea) and can be easily managed [148].

Corticosteroids

Corticosteroids (CSs) have the property of inducing the remission of inflammatory bowel disease through a systemic anti-inflammatory effect. It is mediated by the binding of the steroid molecule to the nuclear receptor, which results in low transcription of genes responsible for the pro-inflammatory proteins and stimulation of genes

that are responsible for the anti-inflammatory effects [155]. In IBD, CSs are recommended for patients who have been receiving mesalazine for more than 4 weeks for moderate forms, and only after the complete cycle of the treatment [156]. The systemic route of corticosteroid administration can cause different effects: diabetes, hypertension, immunosuppression predisposing to infections with opportunistic germs, osteoporosis, etc. [157,158]. In the case of long-term use, corticosteroid dependence may also occur, being defined by the use of high doses of steroids for the same or no effect [159]. The second generation of corticosteroids (butenonide) has a higher safety profile than the others, being administered orally with gradual release in the terminal ileum and ascending colon (the pH-dependent capsules decreasing its systemic effects) [160]. For mild or severe forms of CD, with lesions placed on the ascending colon or the terminal ileum, 9 mg daily therapy for 8 weeks is generally effective [156].

Immunomodulators

Thiopurines

T-lymphocytes are an important element in the pathogenesis of inflammatory bowel disease, penetrating the mucosa and releasing pro-inflammatory cytokines, which further cause local lesions. Thiopurines, such as azathioprine and mercaptopurine, inhibit T-lymphocytes proliferation and activation [161]. In an observational study, thiopurines can maintain the remission of UC for 7 years in 43.8% of the cases, and prevent colectomy in 88% of the cases [162]. The effect of azathioprine is favorable in both UC and CD, reducing the rate of hospitalization and preventing postoperative complications [163-165]. In the case of CD, azathioprine determines the remission of the disease and reduces the need for steroids to no dose for 60 months [166]. However, the side effects of thiopurines are multiple and can be life-threatening. Such adverse effects are represented by the suppression of bone marrow, hepatic cytolysis with hepatitis [167] and gastrointestinal disorders [168]. For this reason, 39% of patients discontinue the treatment with thiopurine within the first 3 months [169].

Methotrexate

At low doses, methotrexate causes the inhibition of DNA synthesis, as well as the lowering of the plasmatic levels of IL-1, IL-2, IL-6, IL-18, all this leading to a decrease in the inflammatory response and the activity of T-lymphocytes [170]. Administration of 25mg methotrexate/week can maintain the remission of CD for 40 weeks, the remission being noted in 72% of the patient after 3 months of treatment [171]. Methotrexate side effects are represented by nausea, vomiting, fatigue, and immunosuppression which can lead to atypical pneumonia. Regarding UC, methotrexate has not been proven its efficacy as an inductor treatment [172].

Calcineurin inhibitors

The calcineurin inhibitors used in the treatment of IBD are cyclosporine A and tacrolimus, which inhibit the NEAT system by binding to specific receptors (Cyclophilin A and FK binding protein 12) [173]. In addition, tacrolimus induces macrophage apoptosis and inhibits the release of pro-inflammatory cytokines IL-12, IL-23 and TNF α [174]. Cyclosporine A is indicated in severe forms of UC refractory to treatment, by administration of 4mg/kg. Remission can be achieved in 80% of the patients over 8 days, relatively similar results being obtained with a dose lower than 2mg/kg [175,176]. Regarding the association of CSs with cyclosporine A in the relapses of patients with CD, no advantages were observed compared to the CSs monotherapy [177].

In comparison with cyclosporine, tacrolimus has a 10-20-time stronger effect in vivo and 30-100 times in vitro [178]. It has the ability of inducing the remission of UC and maintaining it for more than 30 days in 70.4% of the cases [179]. At the same time, it prevents the surgical therapy in the evolution of the patients, with rates of up to 69% at 1 year from the beginning of the treatment [180]. Regarding the effect of tacrolimus in CD, the data are controversial; the remission rates is obtained on about 44.3% of the patients, regardless of the route of administration (topical or systemic) [181]. The side effects of tacrolimus are kidney and liver damages, predisposition to infections and hyperkalemia. The maximum benefit can be obtained by monitoring the plasma level of the drug, which must be maintained at concentrations between 10-15ng/ml [182].

Biologics

Biological treatments target either pro-inflammatory cytokines (TNF α , IL-12, IL-23) or several proteins (integrins). TNF α is a cytokine involved in inflammation, apoptosis and cell proliferation; the overexpression of TNF α gene can lead to various autoimmune diseases [183]. Treatments with anti-TNF α monoclonal antibodies are useful in non-responding patients to immunomodulatory therapy with CSs, or in patients who become dependent on CSs [147]. Infliximab has been shown to be effective; after 54 weeks of treatment, rates of colo-rectomy and hospitalization decrease by 7% for the patients with mild and severe forms of UC [184]. In CD, 5mg/kg of intravenous infliximab led to remission in 68% of the patients, and the fistula cure rate was 55% [185]. Golimumab, a human monoclonal antibody against TNF α , determines a better cure rate than placebo [186,187], so it is currently used in patients with severe forms of UC and non-responding to conventional treatment [188,189]. Although TNF α plays a central role in the pathogenesis of inflammatory bowel disease (its inhibition being very important for successful treatment), there are patients who

do not respond to treatment in 40% of the cases, and between 23-46% of subjects lose their response to treatment over time [190]. This leads to several strategies of improving the therapeutic efficacy, such as de-escalation after remission, or increasing the period between administrations [191].

IL-12 and IL-23 are pro-inflammatory cytokines released in the antigen recognition reaction by the immune cells, being involved in UC and CD in the onset and maintenance of intestinal inflammation [192]. Structurally, IL-12 and IL-23 have two components, p40 that is a common component, and p35 for IL-12 or p19 for IL-23.

Monoclonal antibodies have been developed, with ustekinumab binding to the common component p40. This component prevents cytokine from binding to its own receptor on the surface of T/ NK cells, thus leading to the inhibition of inflammation. Randomized studies have shown the effectiveness of ustekinumab in moderate and severe cases of UC and CD [193]. By administering it for 8 weeks, the endoscopic activity score of the disease decreased by 2.8 points compared to the control group ($p = 0.012$) [194]. Mirikizumab is a monoclonal antibody against the p19 subunit of IL-23, which leads to the clinical remission of the disease by administration of 200 mg for 12 weeks. No major adverse reactions were observed during the administration of these monoclonal antibodies for treatment induction and/or maintenance [195].

Integrins are surface cell receptors that intervene in the leukocytes localization by binding them to specific cellular receptors, called cell adhesion molecules. Integrin $\alpha 4\beta 7$ is present in the intestine and lymphoid tissue, which allows the migration of leukocytes in the mucosa and adhesion to this level. Patients with inflammatory bowel disease have a high number of such receptors [196]. The accumulation of inflammatory cells aggravates local inflammation by the massive release of $\text{INF-}\gamma$, $\text{TNF-}\alpha$, IL-17A, which decrease the tissue repair system [197]. Anti-integrin treatment blocks the effect of integrin on endothelial cells/ leukocytes at the time of contact with the intestinal mucosa.

Vedolizumab is a monoclonal antibody that binds to $\alpha 4\beta 7$, thus preventing the migration of lymphocytes to the intestinal receptor and further reducing the degree of inflammation. Phase 3 studies have shown that vedolizumab can maintain and induce the remission of the disease with a high tolerance for patients, without the risk of immunosuppression. The drug has a high selectivity for receptors [198,199], being indicated for patients with moderate or severe UC/CD that do not respond to conventional or anti-TNF α therapy [200]. If the remission of the disease is not achieved, it can be associated with calcineurin or TNF α inhibitors [201,202].

Etolizumab is an ultra-selective monoclonal antibody that inhibits the $\beta 7$ subunit of the integrin $\alpha 4\beta 7$ and $\alpha E\beta 7$ [197], thus leading to the remission of moderate or severe forms of UC and CD [203].

Biological agents are highly effective, selective and have a low toxicity. Such agents are expensive and the patients may be intolerant or unresponsive to the treatment if used improperly, due to JAK inhibitors that block multiple intercellular or intracellular signaling pathways. However, it can be a solution for unresponsive patients [147]. Tofacitinib is an inhibitor of JAK1 which is found ubiquitously and JAK3 that is found in hematopoietic cells, being useful in the treatment of severe forms of UC [204]. It is able to induce remission after 8 weeks of treatment, the rate of remission in the investigated group being 18.5% higher than in patients in the control group. The best rates of remission of the disease were obtained by administering 10 mg on a daily basis. Long-term side effects are still under investigation; however, thrombosis, infections or the reactivation of varicella-zoster virus may occur in the acute period [205].

Another molecule which was developed and with similar results and uses is filgotinib, which can be administered orally in CD with satisfactory results [206]. If administered in 45 mg doses for moderate/ severe cases of UC, upadacitinib may induce the remission of the disease and improve endoscopic score, but with the risk of developing pulmonary embolism or deep vein thrombosis [207].

Novel Therapies

In refractory or unanswered cases, new methods of inducing the remission of the disease have been developed, including apheresis, cell therapy or exome therapy. They are the starting points for possible new therapeutic methods [147]. Apheresis can reduce the local inflammatory response by isolating and absorbing leukocytes from peripheral blood [208], which can induce the remission of the disease in UC. It has a better efficiency than in the case of CSs administration [209]; in some patients, mucosal healing can be achieved with a low rate of incidence of adverse reactions [210].

Changing the intestinal microbiota

In patients with IBD, there is an imbalance between pathogenic and potentially pathogenic bacteria, which determines an abnormal immune response. Several data suggest that changing the microbiota can improve symptoms and even remit the disease [211]. The antibiotic therapy can be considered a future treatment option, by changing the bacterial flora. Administration may induce some additional beneficial effects, such as the reduction of the risk of postoperative complications [212,213]. In patients with CD, the administration of the antibiotic treatment as the inductor of the remission may lead to modest results, inducing a degree of remission of the disease but without marked symptomatic improvement. Consequently, this class of drugs is not indicated in initiating and maintaining the remission of IBD [214].

Administration of probiotics (living microorganisms-lactobacillus and bifid bacterium), prebiotics and symbiotics have a beneficial effect in IBD, especially in UC. Symbiotics can induce and maintain the remission of the disease better than pro and prebiotics [215], and their association with conventional drugs leads to a superior effect compared to single drug administration [216]. The regular consumption of kefir containing lactobacilli can increase the quality of life of patients with IBD [217]. The effects of probiotics are demonstrated especially in UC. As for CD, the effects of the probiotic therapy are controversial [218].

Fecal material transplant (FMT) is a procedure in which the feces of a healthy donor are transferred to a patient with IBD, in order to improve the intestinal microecology and thus to improve/cure the disease [147]. Fecal transplantation is efficient in the treatment of refractory and recurrent IBD with a success rate of up to 90% [219]. The Australian Consensus approved the transplantation of fecal matter in moderate forms of UC for the first time, in order to induce remission of the disease, the efficiency being superior to placebo after 7 weeks of administration [220]. The main advantage of FMT is the restoration of a quasi-normal ecosystem in patients with IBD, by administration of a wide and varied spectrum of bacteria. The higher the success rate and response to treatment of the disease, the greater the variety of bacteria from the donor [221]. The method of administration of transplant material can influence the efficiency of the treatment. The best results are obtained by administering fecal matter through enemas, being able to administer larger quantities and in a more efficient way (than the administration by superior route or by colonoscopy) [222]. One of the shortcomings of fecal transplantation is the inability to determine long-term efficacy, because studies have not monitored patients for more than 1 year [223]. There may be some side effects such as bloating and diarrhea [224], or other reactions related to the oral route of administration: aspiration pneumonia and the perforation of the colon [225].

Stem cells have the ability of differentiating and proliferating on a particular cell line. They have the ability of repairing damaged tissue in IBD, helping to restore the mucosal barrier. The cells involved in the pathophysiological mechanisms of UC and CD are related to hematopoietic/ mesenchymal line, as well as to intestinal mucosa that can be restored (reducing inflammation and inducing the remission of the disease) [147].

Hematopoietic stem cells have the ability of migrating to damaged tissues, helping them with the restoration and regeneration process. In IBD, the cells used for transplant are harvested from the bone marrow, umbilical cord or peripheral blood with CD34 surface marker [226]. Several studies have been published on this topic. For example, 82 patients with CD received stem cell transplants, which led

to remission and relief of symptoms in 68% of patients, with a disease-free period of 6-174 months [227]. At the same time, stem cell transplants resulted in a 38% reduction in the need for steroids in these patients after 3 months, half of all subjects achieving mucosal healing [228].

Regarding the safety of this new method of treatment, literature data show that patients need maintenance or rescue treatment after 1 year of treatment [229]. About 80% of the patients reach the initial stage of the disease or require a new transplant after 5 years [230]. All this are associated with an increased risk of severe immunosuppression, leading to viral infections, sepsis and pneumonia. For this reason, it is considered that autologous transplantation may be useful in the refractory forms of the disease that do not refer to other therapeutic methods [231].

ABX464 is an antiviral drug that can be used as the only treatment for HIV patients [232]. It may increase the expression of miR-124 and may inhibit the immune response of patients with IBD [233]. By administering 50 mg/ day of ABX464 orally for 8 weeks, an improvement in the endoscopic appearance can be achieved, without severe side effects [234].

IL-10 is a pro-inflammatory cytokine that may be interfered, decreasing UC and CD activity but leading to long-term anemia or thrombocytopenia [235]. AMT101 is a chimeric protein fragment binding to the IL-10 receptor in lamina propria, which appears to stimulate/ exert anti-inflammatory proteins without side effects [236].

Conclusions

From the initial description of inflammatory bowel disease, many pathophysiological mechanisms have been elucidated and presented. These mechanisms can be interfered with by different therapeutic classes of drugs. To note, such mechanisms are induced by several (genetic, environmental, immunological, and microbiotics) factors, which are interdependent and should not be addressed separately. This means that IBD should be considered a multifactorial systemic disease, which must be treated by complex therapeutic methods.

The diagnosis must take into account the digestive manifestations, including their complications, as well as the extradigestive manifestations, which may be difficult for the clinician to detect.

The treatment must be multimodal on all possible levels, by gradually initiating an appropriate therapy. A treatment upgrade may be necessary in the case of non-responsive forms, and even a rescue therapy when life-threatening complications occur.

Finally, a healthy lifestyle without exposure to stress, pollutants, with a rich diet in flavonoids, regular exercise can help the patients relieve their symptoms and induce/ maintain the remission of the disease.

Conflict of interest disclosure

There are no known conflicts of interest in the publication of this article. The manuscript was read and approved by all authors.

Compliance with ethical standards

Any aspect of the work covered in this manuscript has been conducted with the ethical approval of all relevant bodies and that such approvals are acknowledged within the manuscript.

References

- Jostins L, Ripke S, Weersma RK, et al. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature*. 2012 Nov 1; 491(7422):119-24. doi: 10.1038/nature11582
- Inohara N, Ogura Y, Fontalba A, Gutierrez O, Pons F, Crespo J, Fukase K, Inamura S, Kusumoto S, Hashimoto M, Foster SJ, Moran AP, Fernandez-Luna JL, Nuñez G. Host recognition of bacterial muramyl dipeptide mediated through NOD2. Implications for Crohn's disease. *J Biol Chem*. 2003 Feb 21;278(8):5509-12. doi: 10.1074/jbc.C200673200
- Kuballa P, Huett A, Rioux JD, Daly MJ, Xavier RJ. Impaired autophagy of an intracellular pathogen induced by a Crohn's disease associated ATG16L1 variant. *PLoS One*. 2008;3(10):e3391. doi: 10.1371/journal.pone.0003391
- Sergiu MI, Elena MM, Marilena M, Andrada D, Emanuela PR. Microbiota: the missing link in the etiology of inflammatory bowel disease. *J Mind Med Sci*. 2020;7(1):29-33. doi: 10.22543/7674.71.P2933
- Duerr RH, Taylor KD, Brant SR, Rioux JD, Silverberg MS, Daly MJ, Steinhart AH, Abraham C, Regueiro M, Griffiths A, Dassopoulos T, Bitton A, Yang H, Targan S, Datta LW, Kistner EO, Schumm LP, Lee AT, Gregersen PK, Barmada MM, Rotter JI, Nicolae DL, Cho JH. A genome-wide association study identifies IL23R as an inflammatory bowel disease gene. *Science*. 2006 Dec 1;314(5804):1461-3. doi: 10.1126/science.1135245
- Anderson CA, Boucher G, Lees CW, et al. Meta-analysis identifies 29 additional ulcerative colitis risk loci, increasing the number of confirmed associations to 47. *Nat Genet*. 2011 Mar;43(3):246-52. doi: 10.1038/ng.764
- Brand S. Crohn's disease: Th1, Th17 or both? The change of a paradigm: new immunological and genetic insights implicate Th17 cells in the pathogenesis of Crohn's disease. *Gut*. 2009 Aug;58(8):1152-67. doi: 10.1136/gut.2008.163667
- Loftus EV Jr. Clinical epidemiology of inflammatory bowel disease: Incidence, prevalence, and environmental influences. *Gastroenterology*. 2004 May; 126(6):1504-17. doi: 10.1053/j.gastro.2004.01.063
- Cosnes J. Tobacco and IBD: relevance in the understanding of disease mechanisms and clinical practice. *Best Pract Res Clin Gastroenterol*. 2004 Jun; 18(3):481-96. doi: 10.1016/j.bpg.2003.12.003
- Peneş NO, Weber B, Păun SD. Role of genetic polymorphism in nutritional supplementation therapy in personalized medicine. *Rom J Morphol Embryol*. 2017;58(1):53-58.
- Birrenbach T, Böcker U. Inflammatory bowel disease and smoking: a review of epidemiology, pathophysiology, and therapeutic implications. *Inflamm Bowel Dis*. 2004 Nov;10(6):848-59. doi: 10.1097/00054725-200411000-00019
- Ananthakrishnan AN, Higuchi LM, Huang ES, Khalili H, Richter JM, Fuchs CS, Chan AT. Aspirin, nonsteroidal anti-inflammatory drug use, and risk for Crohn disease and ulcerative colitis: a cohort study. *Ann Intern Med*. 2012 Mar 6;156(5):350-9. doi: 10.7326/0003-4819-156-5-201203060-00007
- Shaw SY, Blanchard JF, Bernstein CN. Association between the use of antibiotics in the first year of life and pediatric inflammatory bowel disease. *Am J Gastroenterol*. 2010 Dec;105(12):2687-92. doi: 10.1038/ajg.2010.398
- Thia KT, Loftus EV Jr, Sandborn WJ, Yang SK. An update on the epidemiology of inflammatory bowel disease in Asia. *Am J Gastroenterol*. 2008 Dec; 103(12):3167-82. doi: 10.1111/j.1572-0241.2008.02158.x
- Tan WC, Qiu D, Liam BL, Ng TP, Lee SH, van Eeden SF, D'Yachkova Y, Hogg JC. The human bone marrow response to acute air pollution caused by forest fires. *Am J Respir Crit Care Med*. 2000 Apr;161(4 Pt 1):1213-7. doi: 10.1164/ajrccm.161.4.9904084
- van Eeden SF, Tan WC, Suwa T, Mukae H, Terashima T, Fujii T, Qui D, Vincent R, Hogg JC. Cytokines involved in the systemic inflammatory response induced by exposure to particulate matter air pollutants (PM(10)). *Am J Respir Crit Care Med*. 2001 Sep 1;164(5):826-30. doi: 10.1164/ajrccm.164.5.2010160
- Kaplan GG, Hubbard J, Korzenik J, Sands BE, Panaccione R, Ghosh S, Wheeler AJ, Villeneuve PJ. The inflammatory bowel diseases and ambient air pollution: a novel association. *Am J Gastroenterol*. 2010 Nov;105(11):2412-9. doi: 10.1038/ajg.2010.252
- Vernia F, Valvano M, Longo S, Cesaro N, Viscido A, Latella G. Vitamin D in Inflammatory Bowel Diseases. Mechanisms of Action and Therapeutic Implications. *Nutrients*. 2022;14(2):269. doi: 10.3390/nu14020269
- Prevention and management of osteoporosis. *World Health Organ Tech Rep Ser*. 2003;921:1-164.
- Domazetovic V, Iantomasi T, Bonanomi AG, Stio M. Vitamin D regulates claudin-2 and claudin-4

- expression in active ulcerative colitis by p-Stat-6 and Smad-7 signaling. *Int J Colorectal Dis.* 2020 Jul; 35(7):1231-1242. doi: 10.1007/s00384-020-03576-0
21. Chen SW, Wang PY, Zhu J, Chen GW, Zhang JL, Chen ZY, Zuo S, Liu YC, Pan YS. Protective effect of 1,25-dihydroxyvitamin d3 on lipopolysaccharide-induced intestinal epithelial tight junction injury in caco-2 cell monolayers. *Inflammation.* 2015 Feb;38(1):375-83. doi: 10.1007/s10753-014-0041-9
22. Du J, Chen Y, Shi Y, Liu T, Cao Y, Tang Y, Ge X, Nie H, Zheng C, Li YC. 1,25-Dihydroxyvitamin D Protects Intestinal Epithelial Barrier by Regulating the Myosin Light Chain Kinase Signaling Pathway. *Inflamm Bowel Dis.* 2015 Nov;21(11):2495-506. doi: 10.1097/MIB.0000000000000526
23. Zhang YG, Lu R, Xia Y, Zhou D, Petrof E, Claud EC, Sun J. Lack of Vitamin D Receptor Leads to Hyperfunction of Claudin-2 in Intestinal Inflammatory Responses. *Inflamm Bowel Dis.* 2019 Jan 1;25(1):97-110. doi: 10.1093/ibd/izy292
24. Garg M, Royce SG, Tikellis C, Shallue C, Sluka P, Wardan H, Hosking P, Monagle S, Thomas M, Lubel JS, Gibson PR. The intestinal vitamin D receptor in inflammatory bowel disease: inverse correlation with inflammation but no relationship with circulating vitamin D status. *Therap Adv Gastroenterol.* 2019;12:1756284818822566. doi: 10.1177/1756284818822566
25. Abreu-Delgado Y, Isidro RA, Torres EA, González A, Cruz ML, Isidro AA, González-Keelan CI, Medero P, Appleyard CB. Serum vitamin D and colonic vitamin D receptor in inflammatory bowel disease. *World J Gastroenterol.* 2016 Apr 7;22(13):3581-91. doi: 10.3748/wjg.v22.i13.3581
26. Chen Y, Du J, Zhang Z, Liu T, Shi Y, Ge X, Li YC. MicroRNA-346 mediates tumor necrosis factor α -induced downregulation of gut epithelial vitamin D receptor in inflammatory bowel diseases. *Inflamm Bowel Dis.* 2014 Nov;20(11):1910-8. doi: 10.1097/MIB.0000000000000158
27. Raftery T, Martineau AR, Greiller CL, Ghosh S, McNamara D, Bennett K, Meddings J, O'Sullivan M. Effects of vitamin D supplementation on intestinal permeability, cathelicidin and disease markers in Crohn's disease: Results from a randomised double-blind placebo-controlled study. *United European Gastroenterol J.* 2015 Jun;3(3):294-302. doi: 10.1177/2050640615572176
28. Garg M, Rosella O, Rosella G, Wu Y, Lubel JS, Gibson PR. Evaluation of a 12-week targeted vitamin D supplementation regimen in patients with active inflammatory bowel disease. *Clin Nutr.* 2018; 37(4):1375-1382. doi: 10.1016/j.clnu.2017.06.011
29. Li M, Weigmann B. A Novel Pathway of Flavonoids Protecting against Inflammatory Bowel Disease: Modulating Enteroendocrine System. *Metabolites.* 2022 Jan 1;12(1):31. doi: 10.3390/metabo12010031
30. Alzogaibi MA. Concepts of oxidative stress and antioxidant defense in Crohn's disease. *World J Gastroenterol.* 2013 Oct 21;19(39):6540-7. doi: 10.3748/wjg.v19.i39.6540
31. Veljaca M, Lesch CA, Pillana R, Sanchez B, Chan K, Guglietta A. BPC-15 reduces trinitrobenzene sulfonic acid-induced colonic damage in rats. *J Pharmacol Exp Ther.* 1995 Jan;272(1):417-22.
32. Noda S, Tanabe S, Suzuki T. Differential effects of flavonoids on barrier integrity in human intestinal Caco-2 cells. *J Agric Food Chem.* 2012 May 9;60(18):4628-33. doi: 10.1021/jf300382h
33. Brückner M, Westphal S, Domschke W, Kucharzik T, Lügering A. Green tea polyphenol epigallocatechin-3-gallate shows therapeutic antioxidative effects in a murine model of colitis. *J Crohns Colitis.* 2012 Mar; 6(2):226-35. doi: 10.1016/j.crohns.2011.08.012
34. Guazelli CFS, Fattori V, Ferraz CR, Borghi SM, Casagrande R, Baracat MM, Verri WA Jr. Antioxidant and anti-inflammatory effects of hesperidin methyl chalcone in experimental ulcerative colitis. *Chem Biol Interact.* 2021;333:109315. doi: 10.1016/j.cbi.2020.109315
35. Hong Z, Piao M. Effect of Quercetin Monoglycosides on Oxidative Stress and Gut Microbiota Diversity in Mice with Dextran Sodium Sulphate-Induced Colitis. *Biomed Res Int.* 2018 Nov 12;2018:8343052. doi: 10.1155/2018/8343052
36. Ren J, Yue B, Wang H, Zhang B, Luo X, Yu Z, Zhang J, Ren Y, Mani S, Wang Z, Dou W. Acacetin Ameliorates Experimental Colitis in Mice via Inhibiting Macrophage Inflammatory Response and Regulating the Composition of Gut Microbiota. *Front Physiol.* 2021; 11:577237. doi: 10.3389/fphys.2020.577237
37. Medzhitov R, Janeway C Jr. Innate immunity. *N Engl J Med.* 2000 Aug 3;343(5):338-44. doi: 10.1056/NEJM200008033430506
38. Bonen DK, Ogura Y, Nicolae DL, Inohara N, Saab L, Tanabe T, Chen FF, Foster SJ, Duerr RH, Brant SR, Cho JH, Nuñez G. Crohn's disease-associated NOD2 variants share a signaling defect in response to lipopolysaccharide and peptidoglycan. *Gastroenterology.* 2003;124(1):140-6. doi: 10.1053/gast.2003.50019
39. Wehkamp J, Harder J, Weichenthal M, Schwab M, Schäffeler E, Schlee M, Herrlinger KR, Stallmach A, Noack F, Fritz P, Schröder JM, Bevins CL, Fellermann K, Stange EF. NOD2 (CARD15) mutations in Crohn's disease are associated with diminished mucosal alpha-defensin expression. *Gut.* 2004 Nov;53(11):1658-64. doi: 10.1136/gut.2003.032805
40. Abraham C, Cho JH. Functional consequences of NOD2 (CARD15) mutations. *Inflamm Bowel Dis.* 2006;12(7):641-50. doi: 10.1097/01.MIB.0000225332.83861.5f

41. Watanabe T, Kitani A, Murray PJ, Strober W. NOD2 is a negative regulator of Toll-like receptor 2-mediated T helper type 1 responses. *Nat Immunol.* 2004 Aug; 5(8):800-8. doi: 10.1038/ni1092
42. Korn T, Bettelli E, Oukka M, Kuchroo VK. IL-17 and Th17 Cells. *Annu Rev Immunol.* 2009;27:485-517. doi: 10.1146/annurev.immunol.021908.132710
43. Breese E, Braegger CP, Corrigan CJ, Walker-Smith JA, MacDonald TT. Interleukin-2- and interferon-gamma-secreting T cells in normal and diseased human intestinal mucosa. *Immunology.* 1993 Jan;78(1):127-31.
44. Heller F, Florian P, Bojarski C, Richter J, Christ M, Hillenbrand B, Mankertz J, Gitter AH, Bürgel N, Fromm M, Zeitz M, Fuss I, Strober W, Schulzke JD. Interleukin-13 is the key effector Th2 cytokine in ulcerative colitis that affects epithelial tight junctions, apoptosis, and cell restitution. *Gastroenterology.* 2005 Aug;129(2):550-64. doi: 10.1016/j.gastro.2005.05.002
45. Di Sabatino A, Biancheri P, Rovedatti L, MacDonald TT, Corazza GR. New pathogenic paradigms in inflammatory bowel disease. *Inflamm Bowel Dis.* 2012 Feb;18(2):368-71. doi: 10.1002/ibd.21735
46. Rovedatti L, Kudo T, Biancheri P, Sarra M, Knowles CH, Rampton DS, Corazza GR, Monteleone G, Di Sabatino A, Macdonald TT. Differential regulation of interleukin 17 and interferon gamma production in inflammatory bowel disease. *Gut.* 2009 Dec; 58(12):1629-36. doi: 10.1136/gut.2009.182170
47. Vainer B, Nielsen OH, Hendel J, Horn T, Kirman I. Colonic expression and synthesis of interleukin 13 and interleukin 15 in inflammatory bowel disease. *Cytokine.* 2000;12(10):1531-6. doi: 10.1006/cyto.2000.0744
48. Bernardo D, Vallejo-Díez S, Mann ER, Al-Hassi HO, Martínez-Abad B, Montalvillo E, Tee CT, Murugananthan AU, Núñez H, Peake ST, Hart AL, Fernández-Salazar L, Garrote JA, Arranz E, Knight SC. IL-6 promotes immune responses in human ulcerative colitis and induces a skin-homing phenotype in the dendritic cells and T cells they stimulate. *Eur J Immunol.* 2012 May;42(5):1337-53. doi: 10.1002/eji.201142327
49. Zhou L, Ivanov II, Spolski R, Min R, Shenderov K, Egawa T, Levy DE, Leonard WJ, Littman DR. IL-6 programs T(H)-17 cell differentiation by promoting sequential engagement of the IL-21 and IL-23 pathways. *Nat Immunol.* 2007 Sep;8(9):967-74. doi: 10.1038/ni1488
50. Mekhjian HS, Switz DM, Melnyk CS, Rankin GB, Brooks RK. Clinical features and natural history of Crohn's disease. *Gastroenterology.* 1979 Oct;77(4 Pt 2):898-906.
51. Flynn S, Eisenstein S. Inflammatory Bowel Disease Presentation and Diagnosis. *Surg Clin North Am.* 2019 Dec;99(6):1051-1062. doi: 10.1016/j.suc.2019.08.001
52. Greenstein AJ, Sachar DB, Gibas A, Schrag D, Heimann T, Janowitz HD, Aufses AH Jr. Outcome of toxic dilatation in ulcerative and Crohn's colitis. *J Clin Gastroenterol.* 1985 Apr;7(2):137-43. doi: 10.1097/00004836-198504000-00007
53. Dumitru FA, Micu SI, Popoiag RE, Musat M, Caloian AD, Calu D, Constantin VD, Balan DG, Nitipir C, Enache F. Intestinal dysbiosis – a new treatment target in the prevention of colorectal cancer. *J Mind Med Sci.* 2021;8(2):221-228. doi: 10.22543/7674.82.P221228
54. Tozer PJ, Whelan K, Phillips RK, Hart AL. Etiology of perianal Crohn's disease: role of genetic, microbiological, and immunological factors. *Inflamm Bowel Dis.* 2009 Oct;15(10):1591-8. doi: 10.1002/ibd.21026
55. Beaugerie L, Seksik P, Nion-Larmurier I, Gendre JP, Cosnes J. Predictors of Crohn's disease. *Gastroenterology.* 2006 Mar;130(3):650-6. doi: 10.1053/j.gastro.2005.12.019
56. Mishra N, Parmar KK, Huda T. The comparison between the medical and the surgical management of chronic anal fissures. *J Clin Invest Surg.* 2021;6(1):11-16. doi: 10.25083/2559.5555/6.1.3
57. De Dombal FT, Watts JM, Watkinson G, Goligher JC. Local complications of ulcerative colitis: stricture, pseudopolypoidosis, and carcinoma of colon and rectum. *Br Med J.* 1966 Jun 11;1(5501):1442-7. doi: 10.1136/bmj.1.5501.1442
58. Bernstein CN, Blanchard JF, Rawsthorne P, Yu N. The prevalence of extraintestinal diseases in inflammatory bowel disease: a population-based study. *Am J Gastroenterol.* 2001;96(4):1116-22. doi: 10.1111/j.1572-0241.2001.03756.x
59. Park SK, Wong Z, Park SH, Vu KV, Bang KB, Piyachaturawat P, Myint T, Hilmi I, Park DI. Extraintestinal manifestation of inflammatory bowel disease in Asian patients: A multinational study. *Dig Liver Dis.* 2021;53(2):196-201. doi: 10.1016/j.dld.2020.06.046
60. Biancone L, Mandal A, Yang H, Dasgupta T, Paoluzi AO, Marcheggiano A, Paoluzi P, Pallone F, Das KM. Production of immunoglobulin G and G1 antibodies to cytoskeletal protein by lamina propria cells in ulcerative colitis. *Gastroenterology.* 1995 Jul;109(1):3-12. doi: 10.1016/0016-5085(95)90263-5
61. Manea M, Marcu D, Motofei I, et al. Cardiovascular risk in patients with inflammatory bowel diseases: a review. *Romanian Biotechnological Letters.* 2019; 24(2):366-373. doi: 10.25083/rbl/24.2/366.373
62. Geng X, Biancone L, Dai HH, Lin JJ, Yoshizaki N, Dasgupta A, Pallone F, Das KM. Tropomyosin isoforms in intestinal mucosa: production of autoantibodies to tropomyosin isoforms in ulcerative colitis. *Gastroenterology.* 1998 May;114(5):912-22. doi: 10.1016/s0016-5085(98)70310-5
63. Roussomoustakaki M, Satsangi J, Welsh K, Louis E, Fanning G, Targan S, Landers C, Jewell DP. Genetic

- markers may predict disease behavior in patients with ulcerative colitis. *Gastroenterology*. 1997 Jun;112(6):1845-53. doi: 10.1053/gast.1997.v112.pm9178675
64. Ott C, Schölmerich J. Extraintestinal manifestations and complications in IBD. *Nat Rev Gastroenterol Hepatol*. 2013;10(10):585-95. doi: 10.1038/nrgastro.2013.117
65. Orchard TR, Thiyagaraja S, Welsh KI, Wordsworth BP, Hill Gaston JS, Jewell DP. Clinical phenotype is related to HLA genotype in the peripheral arthropathies of inflammatory bowel disease. *Gastroenterology*. 2000;118(2):274-8. doi: 10.1016/s0016-5085(00)70209-5
66. Smale S, Natt RS, Orchard TR, Russell AS, Bjarnason I. Inflammatory bowel disease and spondylarthropathy. *Arthritis Rheum*. 2001 Dec;44(12):2728-36. doi: 10.1002/1529-0131(200112)44:12<2728::aid-art459>3.0.co;2-8
67. Vavricka SR, Schoepfer A, Scharl M, Lakatos PL, Navarini A, Rogler G. Extraintestinal Manifestations of Inflammatory Bowel Disease. *Inflamm Bowel Dis*. 2015;21(8):1982-92. doi: 10.1097/MIB.0000000000000392
68. Su CG, Judge TA, Lichtenstein GR. Extraintestinal manifestations of inflammatory bowel disease. *Gastroenterol Clin North Am*. 2002 Mar;31(1):307-27. doi: 10.1016/s0889-8553(01)00019-x
69. Rodríguez-Reyna TS, Martínez-Reyes C, Yamamoto-Furusho JK. Rheumatic manifestations of inflammatory bowel disease. *World J Gastroenterol*. 2009 Nov 28;15(44):5517-24. doi: 10.3748/wjg.15.5517
70. Sandborn WJ, Stenson WF, Brynskov J, Lorenz RG, Steidle GM, Robbins JL, Kent JD, Bloom BJ. Safety of celecoxib in patients with ulcerative colitis in remission: a randomized, placebo-controlled, pilot study. *Clin Gastroenterol Hepatol*. 2006 Feb;4(2):203-11. doi: 10.1016/j.cgh.2005.12.002
71. Farmer RG, Hawk WA, Turnbull RB Jr. Clinical patterns in Crohn's disease: a statistical study of 615 cases. *Gastroenterology*. 1975 Apr;68(4 Pt 1):627-35.
72. Fornaciari G, Salvarani C, Beltrami M, Macchioni P, Stockbrügger RW, Russel MG. Musculoskeletal manifestations in inflammatory bowel disease. *Can J Gastroenterol*. 2001 Jun;15(6):399-403. doi: 10.1155/2001/612531. PMID: 11429669.
73. Braun J, Sieper J. The sacroiliac joint in the spondyloarthropathies. *Curr Opin Rheumatol*. 1996 Jul;8(4):275-87. doi: 10.1097/00002281-199607000-00003. PMID: 8864578.
74. Breban M, Gombert B, Amor B, Dougados M. Efficacy of thalidomide in the treatment of refractory ankylosing spondylitis. *Arthritis Rheum*. 1999;42(3):580-1.
75. Karslioglu B, Tekin AC, Tekin E, Tasatan E. Ankle arthrodesis using computer assisted external fixator in patients with end-stage ankle arthrosis. *J Clin Invest Surg*. 2021;6(2):116-120. doi: 10.25083/2559.5555/6.2.5
76. Van den Bosch F, Kruithof E, De Vos M, De Keyser F, Mielants H. Crohn's disease associated with spondyloarthropathy: effect of TNF-alpha blockade with infliximab on articular symptoms. *Lancet*. 2000 Nov 25;356(9244):1821-2. doi: 10.1016/s0140-6736(00)03239-6
77. Vavricka SR, Brun L, Ballabeni P, Pittet V, Prinz Vavricka BM, Zeitz J, Rogler G, Schoepfer AM. Frequency and risk factors for extraintestinal manifestations in the Swiss inflammatory bowel disease cohort. *Am J Gastroenterol*. 2011 Jan;106(1):110-9. doi: 10.1038/ajg.2010.343
78. Greenstein AJ, Janowitz HD, Sachar DB. The extraintestinal complications of Crohn's disease and ulcerative colitis: a study of 700 patients. *Medicine (Baltimore)*. 1976;55(5):401-12. doi: 10.1097/00005792-197609000-00004
79. Timani S, Mutasim DF. Skin manifestations of inflammatory bowel disease. *Clin Dermatol*. 2008;26(3):265-73. doi: 10.1016/j.clindermatol.2007.10.018
80. Kugathasan S, Miranda A, Nocton J, Drolet BA, Raasch C, Binion DG. Dermatologic manifestations of Crohn disease in children: response to infliximab. *J Pediatr Gastroenterol Nutr*. 2003 Aug;37(2):150-4. doi: 10.1097/00005176-200308000-00013
81. Ortego-Centeno N, Callejas-Rubio JL, Sanchez-Cano D, Caballero-Morales T. Refractory chronic erythema nodosum successfully treated with adalimumab. *J Eur Acad Dermatol Venereol*. 2007 Mar;21(3):408-10. doi: 10.1111/j.1468-3083.2006.01893.x
82. Quin A, Kane S, Ulitsky O. A case of fistulizing Crohn's disease and erythema nodosum managed with adalimumab. *Nat Clin Pract Gastroenterol Hepatol*. 2008 May;5(5):278-81. doi: 10.1038/ncpgasthep1099
83. Callen JP. Pyoderma gangrenosum. *Lancet*. 1998;351(9102):581-5. doi: 10.1016/S0140-6736(97)10187-8
84. Orlov-Slavu C, Parosanu A, Olaru M, et al. How opportune is multigene testing in metastatic colorectal cancer? A review. *J Mind Med Sci*. 2021;8(2):215-220. doi: 10.22543/7674.82.P215220
85. Bennett ML, Jackson JM, Jorizzo JL, Fleischer AB Jr, White WL, Callen JP. Pyoderma gangrenosum. A comparison of typical and atypical forms with an emphasis on time to remission. Case review of 86 patients from 2 institutions. *Medicine (Baltimore)*. 2000 Jan;79(1):37-46. doi: 10.1097/00005792-200001000-00004
86. Farhi D, Cosnes J, Zizi N, Chosidow O, Seksik P, Beaugerie L, Aractingi S, Khosrotehrani K. Significance of erythema nodosum and pyoderma gangrenosum in inflammatory bowel diseases: a cohort study of 2402 patients. *Medicine (Baltimore)*. 2008;87(5):281-293. doi: 10.1097/MD.0b013e318187cc9c
87. Powell RJ, Holbrook MR, Stevens A. Pyoderma gangrenosum and its treatment. *Lancet*. 1997;350(9093):1720-1. doi: 10.1016/S0140-6736(05)63568-4

88. Wollina U, Haroske G. Pyoderma gangraenosum. *Curr Opin Rheumatol*. 2011 Jan;23(1):50-6. doi: 10.1097/BOR.0b013e328341152f
89. Motofei IG, Rowland DL, Georgescu SR, Tampa M, Paunica S, Constantin VD, Balalau C, Manea M, Baleanu BC, Sinescu I. Post-Finasteride Adverse Effects in Male Androgenic Alopecia: A Case Report of Vitiligo. *Skin Pharmacol Physiol*. 2017;30(1):42-45. doi: 10.1159/000455972
90. Becuwe C, Delaporte E, Colombel JF, Piette F, Cortot A, Bergoend H. Sweet's syndrome associated with Crohn's disease. *Acta Derm Venereol*. 1989;69(5):444-5.
91. Benton EC, Rutherford D, Hunter JA. Sweet's syndrome and pyoderma gangrenosum associated with ulcerative colitis. *Acta Derm Venereol*. 1985;65(1):77-80.
92. Kemmett D, Hunter JA. Sweet's syndrome: a clinicopathologic review of twenty-nine cases. *J Am Acad Dermatol*. 1990 Sep;23(3 Pt 1):503-7. doi: 10.1016/0190-9622(90)70250-1
93. Vavricka SR, Manser CN, Hediger S, et al. Periodontitis and gingivitis in inflammatory bowel disease: a case-control study. *Inflamm Bowel Dis*. 2013;19(13): 2768-77. doi: 10.1097/01.MIB.0000438356.84263.3b
94. Thrash B, Patel M, Shah KR, Boland CR, Menter A. Cutaneous manifestations of gastrointestinal disease: part II. *J Am Acad Dermatol*. 2013 Feb;68(2):211.e1-33; quiz 244-6. doi: 10.1016/j.jaad.2012.10.036
95. Petrelli EA, McKinley M, Troncale FJ. Ocular manifestations of inflammatory bowel disease. *Ann Ophthalmol*. 1982 Apr;14(4):356-60.
96. Fries W, Giofré MR, Catanoso M, Lo Gullo R. Treatment of acute uveitis associated with Crohn's disease and sacroileitis with infliximab. *Am J Gastroenterol*. 2002; 97(2):499-500. doi: 10.1111/j.1572-0241.2002.05514.x
97. Yarur AJ, Czul F, Levy C. Hepatobiliary manifestations of inflammatory bowel disease. *Inflamm Bowel Dis*. 2014 Sep;20(9):1655-67. doi: 10.1097/MIB.0000000000000065
98. Olsson R, Danielsson A, Järnerot G, Lindström E, Lööf L, Rolny P, Rydén BO, Tysk C, Wallerstedt S. Prevalence of primary sclerosing cholangitis in patients with ulcerative colitis. *Gastroenterology*. 1991 May; 100(5 Pt 1):1319-23.
99. Tischendorf JJ, Hecker H, Krüger M, Manns MP, Meier PN. Characterization, outcome, and prognosis in 273 patients with primary sclerosing cholangitis: A single center study. *Am J Gastroenterol*. 2007;102(1): 107-14. doi: 10.1111/j.1572-0241.2006.00872.x
100. Lindor KD. Ursodiol for primary sclerosing cholangitis. Mayo Primary Sclerosing Cholangitis-Ursodeoxycholic Acid Study Group. *N Engl J Med*. 1997; 336(10):691-5. doi: 10.1056/NEJM199703063361003
101. Bharadwaj S, Tandon P, Kulkarni G, Rivas J, Charles R. The role of endoscopy in inflammatory bowel disease. *J Dig Dis*. 2015 Dec;16(12):689-98. doi: 10.1111/1751-2980.12301
102. Mowat C, Cole A, Windsor A, Ahmad T, Arnott I, Driscoll R, Mitton S, Orchard T, Rutter M, Younge L, Lees C, Ho GT, Satsangi J, Bloom S; IBD Section of the British Society of Gastroenterology. Guidelines for the management of inflammatory bowel disease in adults. *Gut*. 2011 May;60(5):571-607. doi: 10.1136/gut.2010.224154
103. Spiceland CM, Lodhia N. Endoscopy in inflammatory bowel disease: Role in diagnosis, management, and treatment. *World J Gastroenterol*. 2018 Sep 21;24(35): 4014-4020. doi: 10.3748/wjg.v24.i35.4014
104. Waye JD. The role of colonoscopy in the differential diagnosis of inflammatory bowel disease. *Gastrointest Endosc*. 1977 Feb;23(3):150-4. doi: 10.1016/s0016-5107(77)73622-3
105. Pera A, Bellando P, Caldera D, Ponti V, Astegiano M, Barletti C, David E, Arrigoni A, Rocca G, Verme G. Colonoscopy in inflammatory bowel disease. Diagnostic accuracy and proposal of an endoscopic score. *Gastroenterology*. 1987 Jan;92(1):181-5.
106. Iacucci M, Furfaro F, Matsumoto T, Uraoka T, Smith S, Ghosh S, Kiesslich R. Advanced endoscopic techniques in the assessment of inflammatory bowel disease: new technology, new era. *Gut*. 2019 Mar; 68(3):562-572. doi: 10.1136/gutjnl-2017-315235
107. Subramanian V, Mannath J, Ragunath K, Hawkey CJ. Meta-analysis: the diagnostic yield of chromoendoscopy for detecting dysplasia in patients with colonic inflammatory bowel disease. *Aliment Pharmacol Ther*. 2011 Feb;33(3):304-12. doi: 10.1111/j.1365-2036.2010.04525.x
108. Kiesslich R, Fritsch J, Holtmann M, Koehler HH, Stolte M, Kanzler S, Nafe B, Jung M, Galle PR, Neurath MF. Methylene blue-aided chromoendoscopy for the detection of intraepithelial neoplasia and colon cancer in ulcerative colitis. *Gastroenterology*. 2003 Apr; 124(4):880-8. doi: 10.1053/gast.2003.50146
109. Ibarra-Palomino J, Barreto-Zúñiga R, Elizondo-Rivera J, Bobadilla-Díaz J, Villegas-Jiménez A. Aplicación de la cromoendoscopia para evaluar la gravedad y variabilidad interobservador en la colitis ulcerativa crónica inespecífica (CUCI) [Application of chromoendoscopy to evaluate the severity and interobserver variation in chronic non-specific ulcerative colitis]. *Rev Gastroenterol Mex*. 2002 Oct-Dec;67(4):236-40.
110. Neumann H, Neurath MF, Mudter J. New endoscopic approaches in IBD. *World J Gastroenterol*. 2011 Jan 7;17(1):63-8. doi: 10.3748/wjg.v17.i1.63
111. Neumann H, Fujishiro M, Wilcox CM, Mönkemüller K. Present and future perspectives of virtual chromoendoscopy with i-scan and optical enhancement

- technology. *Dig Endosc*. 2014 Jan;26 Suppl 1:43-51. doi: 10.1111/den.12190
112. Kudo T, Matsumoto T, Esaki M, Yao T, Iida M. Mucosal vascular pattern in ulcerative colitis: observations using narrow band imaging colonoscopy with special reference to histologic inflammation. *Int J Colorectal Dis*. 2009 May;24(5):495-501. doi: 10.1007/s00384-008-0631-9
113. Sasanuma S, Ohtsuka K, Kudo SE, Ogata N, Maeda Y, Misawa M, Mori Y, Kudo T, Hisayuki T, Wakamura K, Hayashi T, Katagiri A, Miyachi H, Baba T, Ishida F. Narrow band imaging efficiency in evaluation of mucosal healing/relapse of ulcerative colitis. *Endosc Int Open*. 2018 May;6(5):E518-E523. doi: 10.1055/s-0044-102297
114. Iacucci M, Kiesslich R, Gui X, Panaccione R, Heatherington J, Akinola O, Ghosh S. Beyond white light: optical enhancement in conjunction with magnification colonoscopy for the assessment of mucosal healing in ulcerative colitis. *Endoscopy*. 2017 Jun;49(6):553-559. doi: 10.1055/s-0042-124363
115. Klenske E, Atreya R, Hartmann A, Fischer S, Hirschmann S, Zundler S, Iacucci M, Neurath MF, Rath T. Magnification endoscopy with optical chromoendoscopy shows strong correlation with histologic inflammation in patients with inflammatory bowel disease. *Endosc Int Open*. 2019 Aug;7(8):E1018-E1026. doi: 10.1055/a-0953-1334
116. Neumann H, Vieth M, Neurath MF, Atreya R. Endocytoscopy allows accurate in vivo differentiation of mucosal inflammatory cells in IBD: a pilot study. *Inflamm Bowel Dis*. 2013 Feb;19(2):356-62. doi: 10.1002/ibd.23025
117. Nakazato Y, Naganuma M, Sugimoto S, Bessho R, Arai M, Kiyohara H, Ono K, Nanki K, Mutaguchi M, Mizuno S, Kobayashi T, Hosoe N, Shimoda M, Abe T, Inoue N, Ogata H, Iwao Y, Kanai T. Endocytoscopy can be used to assess histological healing in ulcerative colitis. *Endoscopy*. 2017 Jun;49(6):560-563. doi: 10.1055/s-0043-106556
118. Neumann H, Fuchs FS, Vieth M, Atreya R, Siebler J, Kiesslich R, Neurath MF. Review article: in vivo imaging by endocytoscopy. *Aliment Pharmacol Ther*. 2011 Jun;33(11):1183-93. doi: 10.1111/j.1365-2036.2011.04647.x
119. Macé V, Ahluwalia A, Coron E, Le Rhun M, Boureille A, Bossard C, Mosnier JF, Matysiak-Budnik T, Tarnawski AS. Confocal laser endomicroscopy: a new gold standard for the assessment of mucosal healing in ulcerative colitis. *J Gastroenterol Hepatol*. 2015 Mar; 30 Suppl 1:85-92. doi: 10.1111/jgh.12748
120. Hundorfean G, Chiriact MT, Mihai S, Hartmann A, Mudter J, Neurath MF. Development and Validation of a Confocal Laser Endomicroscopy-Based Score for In Vivo Assessment of Mucosal Healing in Ulcerative Colitis Patients. *Inflamm Bowel Dis*. 2017 Dec 19; 24(1):35-44. doi: 10.1093/ibd/izx012
121. Annunziata ML, Caviglia R, Papparella LG, Cicala M. Upper gastrointestinal involvement of Crohn's disease: a prospective study on the role of upper endoscopy in the diagnostic work-up. *Dig Dis Sci*. 2012 Jun;57(6): 1618-23. doi: 10.1007/s10620-012-2072-0
122. Paerregaard A. What does the IBD patient hide in the upper gastrointestinal tract? *Inflamm Bowel Dis*. 2009 Jul;15(7):1101-4. doi: 10.1002/ibd.20862
123. Nugent FW, Roy MA. Duodenal Crohn's disease: an analysis of 89 cases. *Am J Gastroenterol*. 1989 Mar; 84(3):249-54.
124. Levine A, Koletzko S, Turner D, Escher JC, Cucchiara S, de Ridder L, Kolho KL, Veres G, Russell RK, Paerregaard A, Buderus S, Greer ML, Dias JA, Veereman-Wauters G, Lionetti P, Sladek M, Martin de Carpi J, Staiano A, Ruemmele FM, Wilson DC; European Society of Pediatric Gastroenterology, Hepatology, and Nutrition. ESPGHAN revised porto criteria for the diagnosis of inflammatory bowel disease in children and adolescents. *J Pediatr Gastroenterol Nutr*. 2014 Jun;58(6):795-806. doi: 10.1097/MPG.0000000000000239
125. Maaser C, Sturm A, Vavricka SR, Kucharzik T, Fiorino G, Annese V, Calabrese E, Baumgart DC, Bettenworth D, Borralho Nunes P, Burisch J, Castiglione F, Eliakim R, Ellul P, González-Lama Y, Gordon H, Halligan S, Katsanos K, Kopylov U, Kotze PG, Krustinš E, Laghi A, Limdi JK, Rieder F, Rimola J, Taylor SA, Tolan D, van Rheenen P, Verstockt B, Stoker J; European Crohn's and Colitis Organisation [ECCO] and the European Society of Gastrointestinal and Abdominal Radiology [ESGAR]. ECCO-ESGAR Guideline for Diagnostic Assessment in IBD Part 1: Initial diagnosis, monitoring of known IBD, detection of complications. *J Crohns Colitis*. 2019 Feb 1;13(2):144-164. doi: 10.1093/ecco-jcc/jjy113
126. Hilmi I, Kobayashi T. Capsule endoscopy in inflammatory bowel disease: when and how. *Intest Res*. 2020 Jul;18(3):265-274. doi: 10.5217/ir.2019.09165
127. Efthymiou A, Viazis N, Mantzaris G, Papadimitriou N, Tzourmakliotis D, Raptis S, Karamanolis DG. Does clinical response correlate with mucosal healing in patients with Crohn's disease of the small bowel? A prospective, case-series study using wireless capsule endoscopy. *Inflamm Bowel Dis*. 2008 Nov;14(11): 1542-7. doi: 10.1002/ibd.20509
128. Le Berre C, Sandborn WJ, Aridhi S, Devignes MD, Fournier L, Smail-Tabbone M, Danese S, Peyrin-Biroulet L. Application of Artificial Intelligence to Gastroenterology and Hepatology. *Gastroenterology*. 2020;158(1):76-94.e2. doi: 10.1053/j.gastro.2019.08.058

129. Ozawa T, Ishihara S, Fujishiro M, Saito H, Kumagai Y, Shichijo S, Aoyama K, Tada T. Novel computer-assisted diagnosis system for endoscopic disease activity in patients with ulcerative colitis. *Gastrointest Endosc.* 2019;89(2):416-421.e1. doi: 10.1016/j.gie.2018.10.020
130. Bossuyt P, Nakase H, Vermeire S, de Hertogh G, Eelbode T, Ferrante M, Hasegawa T, Willekens H, Ikemoto Y, Makino T, Bisschops R. Automatic, computer-aided determination of endoscopic and histological inflammation in patients with mild to moderate ulcerative colitis based on red density. *Gut.* 2020 Oct;69(10):1778-1786. doi: 10.1136/gutjnl-2019-320056
131. Biernacka KB, Barańska D, Grzelak P, Czkwianianc E, Szabelska-Zakrzewska K. Up-to-date overview of imaging techniques in the diagnosis and management of inflammatory bowel diseases. *Prz Gastroenterol.* 2019;14(1):19-25. doi: 10.5114/pg.2019.83423
132. Gomollón F, Dignass A, Annese V, Tilg H, Van Assche G, Lindsay JO, Peyrin-Biroulet L, Cullen GJ, Daperno M, Kucharzik T, Rieder F, Almer S, Armuzzi A, Harbord M, Langhorst J, Sans M, Chowers Y, Fiorino G, Juillerat P, Mantzaris GJ, Rizzello F, Vavricka S, Gionchetti P; ECCO. 3rd European Evidence-based Consensus on the Diagnosis and Management of Crohn's Disease 2016: Part 1: Diagnosis and Medical Management. *J Crohns Colitis.* 2017 Jan;11(1):3-25. doi: 10.1093/ecco-jcc/jjw168
133. Lee SS, Kim AY, Yang SK, Chung JW, Kim SY, Park SH, Ha HK. Crohn disease of the small bowel: comparison of CT enterography, MR enterography, and small-bowel follow-through as diagnostic techniques. *Radiology.* 2009 Jun;251(3):751-61. doi: 10.1148/radiol.2513081184
134. Haas K, Rubesova E, Bass D. Role of imaging in the evaluation of inflammatory bowel disease: How much is too much? *World J Radiol.* 2016 Feb 28;8(2):124-31. doi: 10.4329/wjr.v8.i2.124
135. Deepak P, Bruining DH. Radiographical evaluation of ulcerative colitis. *Gastroenterol Rep (Oxf).* 2014 Aug; 2(3):169-77. doi: 10.1093/gastro/gou026
136. Park MJ, Lim JS. Computed tomography enterography for evaluation of inflammatory bowel disease. *Clin Endosc.* 2013;46(4):327-66. doi: 10.5946/ce.2013.46.4.327
137. Panés J, Bouzas R, Chaparro M, García-Sánchez V, Gisbert JP, Martínez de Guereñu B, Mendoza JL, Paredes JM, Quiroga S, Ripollés T, Rimola J. Systematic review: the use of ultrasonography, computed tomography and magnetic resonance imaging for the diagnosis, assessment of activity and abdominal complications of Crohn's disease. *Aliment Pharmacol Ther.* 2011 Jul;34(2):125-45. doi: 10.1111/j.1365-2036.2011.04710.x
138. Andersen K, Vogt C, Blondin D, Beck A, Heinen W, Aurich V, Häussinger D, Mödder U, Cohnen M. Multi-detector CT-colonography in inflammatory bowel disease: prospective analysis of CT-findings to high-resolution video colonoscopy. *Eur J Radiol.* 2006 Apr; 58(1):140-6. doi: 10.1016/j.ejrad.2005.11.004
139. Deepak P, Axelrad JE, Ananthakrishnan AN. The Role of the Radiologist in Determining Disease Severity in Inflammatory Bowel Diseases. *Gastrointest Endosc Clin N Am.* 2019 Jul;29(3):447-470. doi: 10.1016/j.giec.2019.02.006
140. Desmond AN, O'Regan K, Curran C, McWilliams S, Fitzgerald T, Maher MM, Shanahan F. Crohn's disease: factors associated with exposure to high levels of diagnostic radiation. *Gut.* 2008 Nov;57(11):1524-9. doi: 10.1136/gut.2008.151415
141. Jamieson DH, Shipman P, Jacobson K. Magnetic resonance imaging of the perineum in pediatric patients with inflammatory bowel disease. *Can J Gastroenterol.* 2013 Aug;27(8):476-80. doi: 10.1155/2013/624141
142. Gee MS, Harisinghani MG. MRI in patients with inflammatory bowel disease. *J Magn Reson Imaging.* 2011 Mar;33(3):527-34. doi: 10.1002/jmri.22504
143. Horsthuis K, Bipat S, Bennink RJ, Stoker J. Inflammatory bowel disease diagnosed with US, MR, scintigraphy, and CT: meta-analysis of prospective studies. *Radiology.* 2008 Apr;247(1):64-79. doi: 10.1148/radiol.2471070611
144. Oussalah A, Laurent V, Bruot O, Bressenot A, Bigard MA, Régent D, Peyrin-Biroulet L. Diffusion-weighted magnetic resonance without bowel preparation for detecting colonic inflammation in inflammatory bowel disease. *Gut.* 2010 Aug;59(8):1056-65. doi: 10.1136/gut.2009.197665
145. Buisson A, Joubert A, Montoriol PF, Da Ines D, Hordonneau C, Pereira B, Garcier JM, Bommelaer G, Petitcolin V. Diffusion-weighted magnetic resonance imaging for detecting and assessing ileal inflammation in Crohn's disease. *Aliment Pharmacol Ther.* 2013 Mar; 37(5):537-45. doi: 10.1111/apt.12201
146. Hordonneau C, Buisson A, Scanzi J, Goutorbe F, Pereira B, Borderon C, Da Ines D, Montoriol PF, Garcier JM, Boyer L, Bommelaer G, Petitcolin V. Diffusion-weighted magnetic resonance imaging in ileocolonic Crohn's disease: validation of quantitative index of activity. *Am J Gastroenterol.* 2014 Jan; 109(1):89-98. doi: 10.1038/ajg.2013.385
147. Cai Z, Wang S, Li J. Treatment of Inflammatory Bowel Disease: A Comprehensive Review. *Front Med (Lausanne).* 2021 Dec 20;8:765474. doi: 10.3389/fmed.2021.765474
148. Ionescu M, Ionescu P, Suceveanu AP, Stoian AP, Motofei I, Ardeleanu V, Parepa IR. Cardiovascular risk estimation in young patients with ankylosing

- spondylitis: A new model based on a prospective study in Constanta County, Romania. *Exp Ther Med*. 2021 May;21(5):529. doi: 10.3892/etm.2021.9961
149. Oh-Oka K, Kojima Y, Uchida K, Yoda K, Ishimaru K, Nakajima S, Hemmi J, Kano H, Fujii-Kuriyama Y, Katoh R, Ito H, Nakao A. Induction of Colonic Regulatory T Cells by Mesalamine by Activating the Aryl Hydrocarbon Receptor. *Cell Mol Gastroenterol Hepatol*. 2017;4(1):135-151. doi: 10.1016/j.jcmgh.2017.03.010
 150. Murray A, Nguyen TM, Parker CE, Feagan BG, MacDonald JK. Oral 5-aminosalicylic acid for induction of remission in ulcerative colitis. *Cochrane Database Syst Rev*. 2020 Aug 12;8(8):CD000543. doi: 10.1002/14651858.CD000543.pub5
 151. Eaden J, Abrams K, Ekbom A, Jackson E, Mayberry J. Colorectal cancer prevention in ulcerative colitis: a case-control study. *Aliment Pharmacol Ther*. 2000 Feb;14(2):145-53. doi: 10.1046/j.1365-2036.2000.00698.x
 152. Akobeng AK, Zhang D, Gordon M, MacDonald JK. Oral 5-aminosalicylic acid for maintenance of medically-induced remission in Crohn's disease. *Cochrane Database Syst Rev*. 2016;9(9):CD003715. doi: 10.1002/14651858.CD003715.pub3
 153. Hart A, Ng SC, Watkins J, Paridaens K, Edwards JO, Fullarton JR, Sonderegger YLY, Ghatnekar O, Ghosh S. The use of 5-aminosalicylates in Crohn's disease: a retrospective study using the UK Clinical Practice Research Datalink. *Ann Gastroenterol*. 2020 Sep-Oct;33(5):500-507. doi: 10.20524/aog.2020.0521
 154. Coward S, Kuenzig ME, Hazlewood G, Clement F, McBrien K, Holmes R, Panaccione R, Ghosh S, Seow CH, Rezaie A, Kaplan GG. Comparative Effectiveness of Mesalamine, Sulfasalazine, Corticosteroids, and Budesonide for the Induction of Remission in Crohn's Disease: A Bayesian Network Meta-analysis. *Inflamm Bowel Dis*. 2017 Mar;23(3):461-472. doi: 10.1097/MIB.0000000000001023
 155. Hayashi R, Wada H, Ito K, Adcock IM. Effects of glucocorticoids on gene transcription. *Eur J Pharmacol*. 2004 Oct 1;500(1-3):51-62. doi: 10.1016/j.ejphar.2004.07.011
 156. Barrett K, Saxena S, Pollok R. Using corticosteroids appropriately in inflammatory bowel disease: a guide for primary care. *Br J Gen Pract*. 2018 Oct;68(675):497-498. doi: 10.3399/bjgp18X699341
 157. Curkovic I, Egbring M, Kullak-Ublick GA. Risks of inflammatory bowel disease treatment with glucocorticosteroids and aminosalicylates. *Dig Dis*. 2013;31(3-4):368-73. doi: 10.1159/000354699
 158. Motofei IG. Nobel Prize for immune checkpoint inhibitors, understanding the immunological switching between immunosuppression and autoimmunity. *Expert Opin Drug Saf*. 2021 Dec 27:1-14. doi: 10.1080/14740338.2022.2020243
 159. Chhaya V, Saxena S, Cecil E, Subramanian V, Curcin V, Majeed A, Pollok RC. Steroid dependency and trends in prescribing for inflammatory bowel disease - a 20-year national population-based study. *Aliment Pharmacol Ther*. 2016 Sep;44(5):482-94. doi: 10.1111/apt.13700
 160. Travis SP, Danese S, Kupcinskas L, Alexeeva O, D'Haens G, Gibson PR, Moro L, Jones R, Ballard ED, Masure J, Rossini M, Sandborn WJ. Once-daily budesonide MMX in active, mild-to-moderate ulcerative colitis: results from the randomised CORE II study. *Gut*. 2014 Mar;63(3):433-41. doi: 10.1136/gutjnl-2012-304258
 161. Shin JY, Wey M, Umutesi HG, Sun X, Simecka J, Heo J. Thiopurine Prodrugs Mediate Immunosuppressive Effects by Interfering with Rac1 Protein Function. *J Biol Chem*. 2016 Jun 24;291(26):13699-714. doi: 10.1074/jbc.M115.694422
 162. Yamada S, Yoshino T, Matsuura M, Kimura M, Koshikawa Y, Minami N, Toyonaga T, Honzawa Y, Nakase H. Efficacy and Safety of Long-Term Thiopurine Maintenance Treatment in Japanese Patients With Ulcerative Colitis. *Intest Res*. 2015 Jul;13(3):250-8. doi: 10.5217/ir.2015.13.3.250
 163. Aliuş C, Bacalbaşa N, Bălălaşu C. Innovative Device for Indocyanine Green Navigational Surgery. *J Mind Med Sci*. 2020;7(1):40-45. doi: 10.22543/7674.71.P4045
 164. Gisbert JP, Niño P, Cara C, Rodrigo L. Comparative effectiveness of azathioprine in Crohn's disease and ulcerative colitis: prospective, long-term, follow-up study of 394 patients. *Aliment Pharmacol Ther*. 2008 Jul;28(2):228-38. doi: 10.1111/j.1365-2036.2008.03732.x
 165. Pugliese D, Aratari A, Festa S, Ferraro PM, Monterubbianesi R, Guidi L, Scribano ML, Papi C, Armuzzi A. Sustained Clinical Efficacy and Mucosal Healing of Thiopurine Maintenance Treatment in Ulcerative Colitis: A Real-Life Study. *Gastroenterol Res Pract*. 2018 Oct 3;2018:4195968. doi: 10.1155/2018/4195968
 166. Pinto AL, Chebli LA, Ribeiro MS, Pace FH, Moraes JP, do Amaral FJ Jr, Gaburri PD, Meirelles de Souza AF, Chebli JM. Azathioprine therapy in steroid-dependent patients with Crohn disease: results of a 10-year longitudinal follow-up study. *Med Sci Monit*. 2009 May;15(5):PI19-26.
 167. Connell WR, Kamm MA, Ritchie JK, Lennard-Jones JE. Bone marrow toxicity caused by azathioprine in inflammatory bowel disease: 27 years of experience. *Gut*. 1993;34(8):1081-5. doi: 10.1136/gut.34.8.1081
 168. Walker TR, Leichter AM. 6-Thioguanine can cause serious liver injury in inflammatory bowel disease patients. *J Pediatr Gastroenterol Nutr*. 2004 Feb;38(2):232-3. doi: 10.1097/00005176-200402000-00028

169. Jharap B, Seinen ML, de Boer NK, van Ginkel JR, Linskens RK, Kneppelhout JC, Mulder CJ, van Bodegraven AA. Thiopurine therapy in inflammatory bowel disease patients: analyses of two 8-year intercept cohorts. *Inflamm Bowel Dis*. 2010 Sep;16(9):1541-9. doi: 10.1002/ibd.21221
170. van Dieren JM, Kuipers EJ, Samsom JN, Nieuwenhuis EE, van der Woude CJ. Revisiting the immunomodulators tacrolimus, methotrexate, and mycophenolate mofetil: their mechanisms of action and role in the treatment of IBD. *Inflamm Bowel Dis*. 2006 Apr; 12(4):311-27. doi: 10.1097/01.MIB.0000209787.19952.53
171. Feagan BG, Fedorak RN, Irvine EJ, Wild G, Sutherland L, Steinhart AH, Greenberg GR, Koval J, Wong CJ, Hopkins M, Hanauer SB, McDonald JW. A comparison of methotrexate with placebo for the maintenance of remission in Crohn's disease. North American Crohn's Study Group Investigators. *N Engl J Med*. 2000;342(22):1627-32. doi: 10.1056/NEJM200006013422202
172. Chande N, Wang Y, MacDonald JK, McDonald JW. Methotrexate for induction of remission in ulcerative colitis. *Cochrane Database Syst Rev*. 2014;2014(8):CD006618. doi: 10.1002/14651858.CD006618.pub3
173. Naganuma M, Fujii T, Watanabe M. The use of traditional and newer calcineurin inhibitors in inflammatory bowel disease. *J Gastroenterol*. 2011 Feb;46(2):129-37. doi: 10.1007/s00535-010-0352-z
174. Yoshino T, Nakase H, Honzawa Y, Matsumura K, Yamamoto S, Takeda Y, Ueno S, Uza N, Masuda S, Inui K, Chiba T. Immunosuppressive effects of tacrolimus on macrophages ameliorate experimental colitis. *Inflamm Bowel Dis*. 2010 Dec;16(12):2022-33. doi: 10.1002/ibd.21318
175. Lichtiger S, Present DH, Kornbluth A, Gelernt I, Bauer J, Galler G, Michelassi F, Hanauer S. Cyclosporine in severe ulcerative colitis refractory to steroid therapy. *N Engl J Med*. 1994 Jun 30;330(26):1841-5. doi: 10.1056/NEJM199406303302601
176. Van Assche G, D'Haens G, Noman M, Vermeire S, Hiele M, Asnong K, Arts J, D'Hoore A, Penninckx F, Rutgeerts P. Randomized, double-blind comparison of 4 mg/kg versus 2 mg/kg intravenous cyclosporine in severe ulcerative colitis. *Gastroenterology*. 2003 Oct; 125(4):1025-31. doi: 10.1016/s0016-5085(03)01214-9
177. Stange EF, Modigliani R, Peña AS, Wood AJ, Feutren G, Smith PR. European trial of cyclosporine in chronic active Crohn's disease: a 12-month study. The European Study Group. *Gastroenterology*. 1995 Sep; 109(3):774-82. doi: 10.1016/0016-5085(95)90384-4
178. Kino T, Hatanaka H, Miyata S, Inamura N, Nishiyama M, Yajima T, Goto T, Okuhara M, Kohsaka M, Aoki H, et al. FK-506, a novel immunosuppressant isolated from a Streptomyces. II. Immunosuppressive effect of FK-506 in vitro. *J Antibiot (Tokyo)*. 1987 Sep;40(9):1256-65. doi: 10.7164/antibiotics.40.1256
179. Yamamoto S, Nakase H, Mikami S, Inoue S, Yoshino T, Takeda Y, Kasahara K, Ueno S, Uza N, Kitamura H, Tamaki H, Matsuura M, Inui K, Chiba T. Long-term effect of tacrolimus therapy in patients with refractory ulcerative colitis. *Aliment Pharmacol Ther*. 2008; 28(5):589-97. doi: 10.1111/j.1365-2036.2008.03764.x
180. Komaki Y, Komaki F, Ido A, Sakuraba A. Efficacy and Safety of Tacrolimus Therapy for Active Ulcerative Colitis; A Systematic Review and Meta-analysis. *J Crohns Colitis*. 2016 Apr;10(4):484-94. doi: 10.1093/ecco-jcc/jjv221
181. McSharry K, Dalzell AM, Leiper K, El-Matary W. Systematic review: the role of tacrolimus in the management of Crohn's disease. *Aliment Pharmacol Ther*. 2011;34(11-12):1282-94. doi: 10.1111/j.1365-2036.2011.04873.x
182. Ogata H, Matsui T, Nakamura M, Iida M, Takazoe M, Suzuki Y, Hibi T. A randomised dose finding study of oral tacrolimus (FK506) therapy in refractory ulcerative colitis. *Gut*. 2006 Sep;55(9):1255-62. doi: 10.1136/gut.2005.081794
183. Aggarwal BB, Gupta SC, Kim JH. Historical perspectives on tumor necrosis factor and its superfamily: 25 years later, a golden journey. *Blood*. 2012 Jan 19;119(3):651-65. doi: 10.1182/blood-2011-04-325225
184. Sandborn WJ, Rutgeerts P, Feagan BG, Reinisch W, Olson A, Johanns J, Lu J, Horgan K, Rachmilewitz D, Hanauer SB, Lichtenstein GR, de Villiers WJ, Present D, Sands BE, Colombel JF. Colectomy rate comparison after treatment of ulcerative colitis with placebo or infliximab. *Gastroenterology*. 2009 Oct;137(4):1250-60; quiz 1520. doi: 10.1053/j.gastro.2009.06.061
185. Present DH, Rutgeerts P, Targan S, Hanauer SB, Mayer L, van Hogezaand RA, Podolsky DK, Sands BE, Braakman T, DeWoody KL, Schaible TF, van Deventer SJ. Infliximab for the treatment of fistulas in patients with Crohn's disease. *N Engl J Med*. 1999;340(18):1398-405. doi: 10.1056/NEJM199905063401804
186. Sandborn WJ, Feagan BG, Marano C, Zhang H, Strauss R, Johanns J, Adedokun OJ, Guzzo C, Colombel JF, Reinisch W, Gibson PR, Collins J, Järnerot G, Rutgeerts P; PURSUIT-Maintenance Study Group. Subcutaneous golimumab maintains clinical response in patients with moderate-to-severe ulcerative colitis. *Gastroenterology*. 2014 Jan;146(1):96-109.e1. doi: 10.1053/j.gastro.2013.06.010
187. Limdi JK. Golimumab for ulcerative colitis: adding perspective to the pursuit. *Frontline Gastroenterol*. 2018 Jul;9(3):232-233. doi: 10.1136/flgastro-2017-100929
188. Martineau C, Flourié B, Wils P, Vaysse T, Altwegg R, Buisson A, Amiot A, Pineton de Chambrun G, Abitbol V, Fumery M, Hébuterne X, Viennot S, Laharie D,

- Beaugerie L, Nancey S, Sokol H; Goli-Crohn Study Group. Efficacy and safety of golimumab in Crohn's disease: a French national retrospective study. *Aliment Pharmacol Ther.* 2017 Dec;46(11-12):1077-1084. doi: 10.1111/apt.14371
189. Adegbola SO, Sahnun K, Warusavitarnae J, Hart A, Tozer P. Anti-TNF Therapy in Crohn's Disease. *Int J Mol Sci.* 2018 Jul 31;19(8):2244. doi: 10.3390/ijms19082244
190. Ben-Horin S, Kopylov U, Chowers Y. Optimizing anti-TNF treatments in inflammatory bowel disease. *Autoimmun Rev.* 2014 Jan;13(1):24-30. doi: 10.1016/j.autrev.2013.06.002
191. Yamada S, Yoshino T, Matsuura M, Minami N, Toyonaga T, Honzawa Y, Tsuji Y, Nakase H. Long-term efficacy of infliximab for refractory ulcerative colitis: results from a single center experience. *BMC Gastroenterol.* 2014 Apr 23;14:80. doi: 10.1186/1471-230X-14-80
192. Neurath MF. IL-23: a master regulator in Crohn disease. *Nat Med.* 2007 Jan;13(1):26-8. doi: 10.1038/nm0107-26
193. Almradi A, Hanzel J, Sedano R, Parker CE, Feagan BG, Ma C, Jairath V. Clinical Trials of IL-12/IL-23 Inhibitors in Inflammatory Bowel Disease. *BioDrugs.* 2020 Dec;34(6):713-721. doi: 10.1007/s40259-020-00451-w
194. Rutgeerts P, Gasink C, Chan D, Lang Y, Pollack P, Colombel JF, Wolf DC, Jacobstein D, Johanns J, Szapary P, Adedokun OJ, Feagan BG, Sandborn WJ. Efficacy of Ustekinumab for Inducing Endoscopic Healing in Patients With Crohn's Disease. *Gastroenterology.* 2018 Oct;155(4):1045-1058. doi: 10.1053/j.gastro.2018.06.035
195. Sandborn WJ, Ferrante M, Bhandari BR, Berliba E, Feagan BG, Hibi T, Tuttle JL, Klekotka P, Friedrich S, Durante M, Morgan-Cox M, Laskowski J, Schmitz J, D'Haens GR. Efficacy and Safety of Mirikizumab in a Randomized Phase 2 Study of Patients With Ulcerative Colitis. *Gastroenterology.* 2020;158(3):537-549.e10. doi: 10.1053/j.gastro.2019.08.043
196. Wyant T, Fedyk E, Abhyankar B. An Overview of the Mechanism of Action of the Monoclonal Antibody Vedolizumab. *J Crohns Colitis.* 2016 Dec;10(12):1437-1444. doi: 10.1093/ecco-jcc/jjw092
197. Tang MT, Keir ME, Erickson R, Stefanich EG, Fuh FK, Ramirez-Montagut T, McBride JM, Danilenko DM. Review article: nonclinical and clinical pharmacology, pharmacokinetics and pharmacodynamics of etrolizumab, an anti- β 7 integrin therapy for inflammatory bowel disease. *Aliment Pharmacol Ther.* 2018 Jun;47(11):1440-1452. doi: 10.1111/apt.14631
198. Feagan BG, Rutgeerts P, Sands BE, Hanauer S, Colombel JF, Sandborn WJ, Van Assche G, Axler J, Kim HJ, Danese S, Fox I, Milch C, Sankoh S, Wyant T, Xu J, Parikh A; GEMINI 1 Study Group. Vedolizumab as induction and maintenance therapy for ulcerative colitis. *N Engl J Med.* 2013 Aug 22;369(8):699-710. doi: 10.1056/NEJMoa1215734
199. Sandborn WJ, Feagan BG, Rutgeerts P, Hanauer S, Colombel JF, Sands BE, Lukas M, Fedorak RN, Lee S, Bressler B, Fox I, Rosario M, Sankoh S, Xu J, Stephens K, Milch C, Parikh A; GEMINI 2 Study Group. Vedolizumab as induction and maintenance therapy for Crohn's disease. *N Engl J Med.* 2013 Aug 22;369(8):711-21. doi: 10.1056/NEJMoa1215739
200. Scribano ML. Vedolizumab for inflammatory bowel disease: From randomized controlled trials to real-life evidence. *World J Gastroenterol.* 2018 Jun 21;24(23):2457-2467. doi: 10.3748/wjg.v24.i23.2457
201. Christensen B, Gibson PR, Micic D, Colman RJ, Goepfing SR, Kassim O, Yarur A, Weber CR, Cohen RD, Rubin DT. Safety and Efficacy of Combination Treatment With Calcineurin Inhibitors and Vedolizumab in Patients With Refractory Inflammatory Bowel Disease. *Clin Gastroenterol Hepatol.* 2019 Feb;17(3):486-493. doi: 10.1016/j.cgh.2018.04.060
202. Pellet G, Stefanescu C, Carbonnel F, Peyrin-Biroulet L, Roblin X, Allimant C, Nachury M, Nancey S, Filippi J, Altwegg R, Brixi H, Fotsing G, de Rosamel L, Shili S, Laharie D; Groupe d'Etude Thérapeutique des Affections Inflammatoires du tube Digestif. Efficacy and Safety of Induction Therapy With Calcineurin Inhibitors in Combination With Vedolizumab in Patients With Refractory Ulcerative Colitis. *Clin Gastroenterol Hepatol.* 2019 Feb;17(3):494-501. doi: 10.1016/j.cgh.2018.08.081
203. Vermeire S, O'Byrne S, Keir M, Williams M, Lu TT, Mansfield JC, Lamb CA, Feagan BG, Panes J, Salas A, Baumgart DC, Schreiber S, Dotan I, Sandborn WJ, Tew GW, Luca D, Tang MT, Diehl L, Eastham-Anderson J, De Hertogh G, Perrier C, Egen JG, Kirby JA, van Assche G, Rutgeerts P. Etrolizumab as induction therapy for ulcerative colitis: a randomised, controlled, phase 2 trial. *Lancet.* 2014 Jul 26;384(9940):309-18. doi: 10.1016/S0140-6736(14)60661-9
204. Tran V, Shammas RM, Sauk JS, Padua D. Evaluating tofacitinib citrate in the treatment of moderate-to-severe active ulcerative colitis: design, development and positioning of therapy. *Clin Exp Gastroenterol.* 2019 May 2;12:179-191. doi: 10.2147/CEG.S150908
205. Sandborn WJ, Su C, Sands BE, D'Haens GR, Vermeire S, Schreiber S, Danese S, Feagan BG, Reinisch W, Niezychowski W, Friedman G, Lawendy N, Yu D, Woodworth D, Mukherjee A, Zhang H, Healey P, Panés J; OCTAVE Induction 1, OCTAVE Induction 2, and OCTAVE Sustain Investigators. Tofacitinib as Induction and Maintenance Therapy for Ulcerative

- Colitis. *N Engl J Med*. 2017 May 4;376(18):1723-1736. doi: 10.1056/NEJMoa1606910
206. Vermeire S, Schreiber S, Petryka R, Kuehbacher T, Hebuterne X, Roblin X, Klopocka M, Goldis A, Wisniewska-Jarosinska M, Baranovsky A, Sike R, Stoyanova K, Tasset C, Van der Aa A, Harrison P. Clinical remission in patients with moderate-to-severe Crohn's disease treated with filgotinib (the FITZROY study): results from a phase 2, double-blind, randomised, placebo-controlled trial. *Lancet*. 2017;389(10066):266-275. doi: 10.1016/S0140-6736(16)32537-5
207. Sandborn WJ, Ghosh S, Panes J, Schreiber S, D'Haens G, Tanida S, Siffledeen J, Enejosa J, Zhou W, Othman AA, Huang B, Higgins PDR. Efficacy of Upadacitinib in a Randomized Trial of Patients With Active Ulcerative Colitis. *Gastroenterology*. 2020 Jun;158(8):2139-2149.e14. doi: 10.1053/j.gastro.2020.02.030
208. Naganuma M, Yokoyama Y, Motoya S, Watanabe K, Sawada K, Hirai F, Yamamoto T, Hanai H, Omori T, Kanai T, Hibi T; CAPTAIN study Group. Efficacy of apheresis as maintenance therapy for patients with ulcerative colitis in an open-label prospective multicenter randomised controlled trial. *J Gastroenterol*. 2020 Apr;55(4):390-400. doi: 10.1007/s00535-019-01651-0
209. Yoshino T, Nakase H, Minami N, Yamada S, Matsuura M, Yazumi S, Chiba T. Efficacy and safety of granulocyte and monocyte adsorption apheresis for ulcerative colitis: a meta-analysis. *Dig Liver Dis*. 2014 Mar;46(3):219-26. doi: 10.1016/j.dld.2013.10.011
210. Fukuchi T, Nakase H, Ubukata S, Matsuura M, Yoshino T, Toyonaga T, Shimazu K, Koga H, Yamashita H, Ito D, Ashida K. Therapeutic effect of intensive granulocyte and monocyte adsorption apheresis combined with thiopurines for steroid- and biologics-naïve Japanese patients with early-diagnosed Crohn's disease. *BMC Gastroenterol*. 2014 Jul 11;13:124. doi: 10.1186/1471-230X-14-124
211. Ocansey DKW, Wang L, Wang J, Yan Y, Qian H, Zhang X, Xu W, Mao F. Mesenchymal stem cell-gut microbiota interaction in the repair of inflammatory bowel disease: an enhanced therapeutic effect. *Clin Transl Med*. 2019 Dec 23;8(1):31. doi: 10.1186/s40169-019-0251-8
212. Khan KJ, Ullman TA, Ford AC, Abreu MT, Abadir A, Marshall JK, Talley NJ, Moayyedi P. Antibiotic therapy in inflammatory bowel disease: a systematic review and meta-analysis. *Am J Gastroenterol*. 2011 Apr;106(4):661-73. doi: 10.1038/ajg.2011.72
213. Uchino M, Ikeuchi H, Bando T, Chohno T, Sasaki H, Horio Y, Nakajima K, Takesue Y. Efficacy of Preoperative Oral Antibiotic Prophylaxis for the Prevention of Surgical Site Infections in Patients With Crohn Disease: A Randomized Controlled Trial. *Ann Surg*. 2019 Mar;269(3):420-426. doi: 10.1097/SLA.0000000000002567
214. Townsend CM, Parker CE, MacDonald JK, Nguyen TM, Jairath V, Feagan BG, Khanna R. Antibiotics for induction and maintenance of remission in Crohn's disease. *Cochrane Database Syst Rev*. 2019;2(2):CD012730. doi: 10.1002/14651858.CD012730.pub2
215. Zhang XF, Guan XX, Tang YJ, Sun JF, Wang XK, Wang WD, Fan JM. Clinical effects and gut microbiota changes of using probiotics, prebiotics or synbiotics in inflammatory bowel disease: a systematic review and meta-analysis. *Eur J Nutr*. 2021 Aug;60(5):2855-2875. doi: 10.1007/s00394-021-02503-5
216. Mardini HE, Grigorian AY. Probiotic mix VSL#3 is effective adjunctive therapy for mild to moderately active ulcerative colitis: a meta-analysis. *Inflamm Bowel Dis*. 2014;20(9):1562-7. doi: 10.1097/MIB.0000000000000084
217. Yılmaz İ, Dolar ME, Özpınar H. Effect of administering kefir on the changes in fecal microbiota and symptoms of inflammatory bowel disease: A randomized controlled trial. *Turk J Gastroenterol*. 2019 Mar;30(3):242-253. doi: 10.5152/tjg.2018.18227
218. Kaur L, Gordon M, Baines PA, Iheozor-Ejiofor Z, Sinopoulou V, Akobeng AK. Probiotics for induction of remission in ulcerative colitis. *Cochrane Database Syst Rev*. 2020;3(3):CD005573. doi: 10.1002/14651858.CD005573.pub3
219. Dumitriu B, Valcea S, Andrei G, Beuran M. Evaluation of anemia as a postoperative risk factor in the evolution of patients with gastric resection for malignancies. *J Clin Invest Surg*. 2021;6(2):136-140. doi: 10.25083/2559.5555/6.2.8
220. Haifer C, Kelly CR, Paramsothy S, Andresen D, Papanicolaos LE, McKew GL, Borody TJ, Kamm M, Costello SP, Andrews JM, Begun J, Chan HT, Connor S, Ghaly S, Johnson PD, Lemberg DA, Paramsothy R, Redmond A, Sheorey H, van der Poorten D, Leong RW. Australian consensus statements for the regulation, production and use of faecal microbiota transplantation in clinical practice. *Gut*. 2020 May;69(5):801-810. doi: 10.1136/gutjnl-2019-320260
221. Stoian AP, Hainarosie R, Pietrosanu C, Rusescu A, Andronache LF, et al. Modern concepts in non-surgical esthetics; a review. *J Mind Med Sci*. 2019;6(2):190-195. doi: 10.22543/7674.62.P190195
222. Paramsothy S, Paramsothy R, Rubin DT, Kamm MA, Kaakoush NO, Mitchell HM, Castaño-Rodríguez N. Faecal Microbiota Transplantation for Inflammatory Bowel Disease: A Systematic Review and Meta-analysis. *J Crohns Colitis*. 2017 Oct 1;11(10):1180-1199. doi: 10.1093/ecco-jcc/jjx063
223. Moayyedi P, Surette MG, Kim PT, Libertucci J, Wolfe M, Onischi C, Armstrong D, Marshall JK, Kassam Z, Reinisch W, Lee CH. Fecal Microbiota Transplantation Induces Remission in Patients With Active Ulcerative Colitis in a Randomized Controlled Trial.

- Gastroenterology*. 2015 Jul;149(1):102-109.e6. doi: 10.1053/j.gastro.2015.04.001
224. Keskin A, Kararlioglu B. Did Covid-19 pandemic narrow the spectrum of surgical indications? *J Clin Invest Surg*. 2021;6(1):58-63. doi: 10.25083/2559.5555/6.1.11
225. Vermeire S, Joossens M, Verbeke K, Wang J, Machiels K, Sabino J, Ferrante M, Van Assche G, Rutgeerts P, Raes J. Donor Species Richness Determines Faecal Microbiota Transplantation Success in Inflammatory Bowel Disease. *J Crohns Colitis*. 2016 Apr;10(4):387-94. doi: 10.1093/ecco-jcc/jjv203
226. Nagaishi K, Arimura Y, Fujimiya M. Stem cell therapy for inflammatory bowel disease. *J Gastroenterol*. 2015 Mar;50(3):280-6. doi: 10.1007/s00535-015-1040-9
227. Brierley CK, Castilla-Llorente C, Labopin M, Badoglio M, Rovira M, Ricart E, Dierickx D, Vermeire S, Hasselblatt P, Finke J, Onida F, Cassinotti A, Satsangi J, Kazmi M, López-Sanromán A, Schmidt C, Farge D, Travis SPL, Hawkey CJ, Snowden JA; European Society for Blood and Marrow Transplantation [EBMT] Autoimmune Diseases Working Party [ADWP]. Autologous Haematopoietic Stem Cell Transplantation for Crohn's Disease: A Retrospective Survey of Long-term Outcomes From the European Society for Blood and Marrow Transplantation. *J Crohns Colitis*. 2018 Aug 29;12(9):1097-1103. doi: 10.1093/ecco-jcc/jjy069
228. Lindsay JO, Allez M, Clark M, Labopin M, Ricart E, Rogler G, Rovira M, Satsangi J, Farge D, Hawkey CJ; ASTIC trial group; European Society for Blood and Marrow Transplantation Autoimmune Disease Working Party; European Crohn's and Colitis Organisation. Autologous stem-cell transplantation in treatment-refractory Crohn's disease: an analysis of pooled data from the ASTIC trial. *Lancet Gastroenterol Hepatol*. 2017 Jun;2(6):399-406. doi: 10.1016/S2468-1253(17)30056-0
229. Snowden JA, Ansari A, Sachchithanatham S, Jackson G, Thompson N, Lobo A, Sanderson J, Kazmi M. Autologous stem cell transplantation in severe treatment-resistant Crohn's disease: long-term follow-up of UK patients treated on compassionate basis. *QJM*. 2014;107(11):871-7. doi: 10.1093/qjmed/hcu095
230. López-García A, Rovira M, Jauregui-Amezaga A, Marín P, Barastegui R, Salas A, Ribas V, Feu F, Elizalde JI, Fernández-Avilés F, Martínez C, Gutiérrez G, Rosiñol L, Carreras E, Urbano A, Lozano M, Cid J, Suárez-Lledó M, Masamunt MC, Comas D, Giner A, Gallego M, Alfaro I, Ordás I, Panés J, Ricart E. Autologous Haematopoietic Stem Cell Transplantation for Refractory Crohn's Disease: Efficacy in a Single-Centre Cohort. *J Crohns Colitis*. 2017 Oct 1;11(10):1161-1168. doi: 10.1093/ecco-jcc/jjx054
231. Mishra R, Dhawan P, Srivastava AS, Singh AB. Inflammatory bowel disease: Therapeutic limitations and prospective of the stem cell therapy. *World J Stem Cells*. 2020 Oct 26;12(10):1050-1066. doi: 10.4252/wjsc.v12.i10.1050
232. Vautrin A, Manchon L, Garcel A, Campos N, Lapasset L, Laaref AM, Bruno R, Gislard M, Dubois E, Scherrer D, Ehrlich JH, Tazi J. Both anti-inflammatory and antiviral properties of novel drug candidate ABX464 are mediated by modulation of RNA splicing. *Sci Rep*. 2019 Jan 28;9(1):792. doi: 10.1038/s41598-018-37813-y
233. Begon-Pescia C, Mielle J, Campose N, Chebli K, Manchon L, Santo J, Apolit C, Martin K, Lapasset L, Vautrin A, Scherrer D, Garcel A, Tazi J, Daien C. THU0199 ABX464, a novel drug in the field of inflammation, increases MIR-124 and modulates macrophages and T-cell functions. *Ann Rheum Dis*. 2020;79(Suppl 1):321-322. doi: 10.1136/annrheumdis-2020-eular.4959
234. Uysal E, Dokur M, Maralcan G. Abdominal surgery in autoimmune and autoimmune related diseases; A review. *J Clin Invest Surg*. 2021;6(2):76-87. doi: 10.25083/2559.5555/6.2.1
235. Vermeire S, Hébuterne X, Napora P, Wisniewska-Jarosinska M, Kiss G, Bourreille A, Przemysław Z, Nitcheu J, Gineste P, Steens JM, Ehrlich H. OP21 ABX464 is safe and efficacious in a proof-of-concept study in ulcerative colitis patients. *Journal of Crohn's and Colitis*. 2019;13(Supplement_1):S014-S015. doi: 10.1093/ecco-jcc/jjy222.020
236. Fay NC, Muthusamy BP, Nyugen LP, Desai RC, Taverner A, MacKay J, Seung M, Hunter T, Liu K, Chandalia A, Coyle MP, Kim HL, Postlethwaite S, Mangat K, Song L, Seto E, Alam A, Olson CV, Feng W, Saberi M, Mahmood TA, Mrsny RJ. A Novel Fusion of IL-10 Engineered to Traffic across Intestinal Epithelium to Treat Colitis. *J Immunol*. 2020 Dec 1; 205(11):3191-3204. doi: 10.4049/jimmunol.2000848