

June 1982

## *Triepeolus Mesillae* (Hymenoptera: Apoidea) in Illinois: A Cleptoparasitic Bee New to the Eastern United States

John K. Bouseman  
*Illinois Natural History Survey*

Follow this and additional works at: <http://scholar.valpo.edu/tgle>

 Part of the [Entomology Commons](#)

---

### Recommended Citation

Bouseman, John K. (1982) "*Triepeolus Mesillae* (Hymenoptera: Apoidea) in Illinois: A Cleptoparasitic Bee New to the Eastern United States," *The Great Lakes Entomologist*: Vol. 15 : No. 2 , Article 11.  
Available at: <http://scholar.valpo.edu/tgle/vol15/iss2/11>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

*TRIEPEOLUS MESILLAE* (HYMENOPTERA: APOIDEA) IN ILLINOIS: A  
CLEPTOPARASITIC BEE NEW TO THE EASTERN UNITED STATES

Cockerell (Ann. Mag. Nat. Hist., Ser. 7, 13:33-42, 1904) named *Triepeolus mesillae* on the basis of a series of specimens which he collected at Mesilla, New Mexico. A second locality for the species was added by Rozen (Amer. Mus. Novitates 2244, 1966) when he described the larva of *T. mesillae* from cells of *Nomia (Epinomia) triangulifera* Vachal excavated by E. A. Cross at Lakeview, Douglas County, Kansas, in 1955. Aside from Cockerell's inclusion of *T. mesillae* in his key to the *Triepeolus* of Colorado (Cockerell, Univ. Colorado Stud. 16:99-126, 1928), no other information concerning the species has been published. The purpose of this paper is to record *T. mesillae* from Illinois and thus to provide a third locality for the species and the first from east of the Mississippi River.

On 29 August 1981, an associate of the author, Tamara F. Bouseman, took a male and a female of *Triepeolus mesillae* from flowers of *Helianthus occidentalis* Riddell in Henderson County, Illinois. The bees were collected between 1300 and 1400 CDST from a population of *H. occidentalis* growing in a sand prairie about 3.5 miles N of Oquawka. During the time of the collection of the two specimens of *T. mesillae*, the author collected a pollen-laden female of the recorded host, *Nomia triangulifera*, from nearby flowers of *Helianthus rigidus* (Cass.) Desf., and other specimens of *N. triangulifera* were taken from sunflowers later on that day (Bouseman, J. Kansas Entomol. Soc., 55:95-96, 1982).

ACKNOWLEDGMENTS: Sincere thanks are due to Paul D. Hurd, Jr., National Museum of Natural History, Smithsonian Institution, Washington, D.C., for his confirmation of the identity of the specimens of *T. mesillae* cited herein, and to Tamara F. Bouseman for her aid in the field. This publication is a contribution of the Illinois Natural History Survey and the Illinois Agricultural Experiment Station, College of Agriculture, University of Illinois at Urbana-Champaign. It was supported in part by Hatch Project No. 12-0361, Biosystematics of Insects.

John K. Bouseman  
Illinois Natural History Survey  
Champaign, IL 61820