

October 1986

## Records of *Dufourea Maura* (Hymenoptera: Halictidae) From Isle Royale National Park, Michigan

Michael S. Arduser  
*University of Missouri*

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

---

### Recommended Citation

Arduser, Michael S. 1986. "Records of *Dufourea Maura* (Hymenoptera: Halictidae) From Isle Royale National Park, Michigan," *The Great Lakes Entomologist*, vol 19 (3)

DOI: <https://doi.org/10.22543/0090-0222.1574>

Available at: <https://scholar.valpo.edu/tgle/vol19/iss3/6>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

**RECORDS OF *DUFOUREA MAURA* (HYMENOPTERA: HALICTIDAE) FROM ISLE ROYALE NATIONAL PARK, MICHIGAN**

Michael S. Arduser<sup>1</sup>

ABSTRACT

*Dufourea maura* is recorded from Isle Royale National Park, Michigan. This is the first report of this western North American species east of the Great Plains states.

---

Recent examinations of specimens in the Entomological Research Collection of Michigan State University and The University of Michigan Museum of Zoology, uncovered 31 specimens of *Dufourea maura* (Cresson) from Isle Royale National Park, Michigan (Keweenaw County). According to Hurd (1979) and George E. Bohart (pers. comm.), *D. maura* is primarily a species of the western United States, occurring as far east as North Dakota and Nebraska. The Isle Royale collections include 27 males and four females, taken between 2–27 July in various years. One male was collected from flowers of *Campanula rotundifolia* L.

No biological data have been published for *D. maura* other than Stevens' note (1919) that the species is a "fairly common bee at flowers of *Campanula rotundifolia*" in North Dakota. Given the oligolectic nature of other species in the genus (Bohart, pers. comm.), it seems possible that *D. maura* may restrict its pollen collecting to the flowers of *Campanula rotundifolia* and related species. *C. rotundifolia* is a widely distributed species that is relatively common on Isle Royale, found on moist, exposed, sandy or rocky substrates.

These appear to be the only records of *Dufourea maura* in the eastern United States. Despite much collecting on the mainland of the Upper Peninsula of Michigan in the past several years by the author and others, no additional collections of this species have been made.

ACKNOWLEDGMENTS

I would like to thank Roland L. Fischer, Michigan State University Entomological Museum, and Mark F. O'Brien, The University of Michigan Museum of Zoology, for permitting me to examine some of the specimens in their care. Thanks are due George E. Bohart, USDA-ARS Bee Biology and Systematics Lab, Logan, Utah, who confirmed the identity of *D. maura*, and John K. Bouseman, James H. Hunt, Mark F. O'Brien, and George E. Bohart for reading the manuscript.

---

<sup>1</sup>Department of Biology, University of Missouri-St. Louis, St. Louis, MO 63121.

LITERATURE CITED

- Hurd, P. D., Jr. 1979. Superfamily Apoidea. pp. 1741–2209 in K.V. Krombein, P.D. Hurd, Jr. D.R. Smith, and B.D. Burks (eds.). Catalog of Hymenoptera in America north of Mexico, Vol. 2 (Aculeata). Smithsonian Inst. Press, Washington.
- Stevens, O. A. 1919. The panurgine bees of North Dakota and a new *Epeolus*. Canadian Entomol. 51:205–210.