

October 1987

## Bibliography of Sequential Sampling Plans in Insect Pest Management Based on Wald's Sequential Probability Ratio Test

Gary W. Fowler  
*University of Michigan*

Ann M. Lynch  
*University of Arizona*

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

---

### Recommended Citation

Fowler, Gary W. and Lynch, Ann M. 1987. "Bibliography of Sequential Sampling Plans in Insect Pest Management Based on Wald's Sequential Probability Ratio Test," *The Great Lakes Entomologist*, vol 20 (3)  
DOI: <https://doi.org/10.22543/0090-0222.1617>  
Available at: <https://scholar.valpo.edu/tgle/vol20/iss3/12>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

**BIBLIOGRAPHY OF SEQUENTIAL SAMPLING PLANS IN INSECT PEST MANAGEMENT BASED ON WALD'S SEQUENTIAL PROBABILITY RATIO TEST**Gary W. Fowler<sup>1</sup> and Ann M. Lynch<sup>2</sup>**ABSTRACT**

This paper contains 65 references dealing with the development of sequential sampling plans in insect pest management based on Wald's Sequential Probability Ratio Test (SPRT), 25 in forest entomology and 40 in agriculture entomology. The insect(s) sampled, whether the decision procedure was based on one or two SPRTs, and the mathematical distribution and probabilities of Type I ( $\alpha$ ) and Type II ( $\beta$ ) errors used to develop the SPRTs are also given for each sequential sampling plan.

---

Sequential analysis includes all of those statistical procedures in which the sample size is not fixed prior to sampling. In sequential hypothesis testing, the number of observations taken in a given sample depends on the conclusiveness of evidence collected, observation by observation, for or against the null hypothesis or standard being tested. In other words, the final pattern and number of observations are not determined prior to sampling. The problem is to test the simple null hypothesis against some simple alternative hypothesis. The sample size is a random variable and is based on some stopping rule.

Test procedures based on sequential techniques require, on the average, a much smaller sample size, i.e., only 40–60% as many observations, than is required by equally reliable procedures based on fixed sample size techniques. When observations are expensive, time consuming, or destructive, sequential procedures seem to have a distinct advantage. Sequential hypothesis testing can be used to classify populations or to accept or reject a specific standard. They are especially useful in surveys.

Much of the theoretical literature in sequential hypothesis testing deals with Wald's Sequential Probability Ratio Test (SPRT) (Ghosh 1970; Wald 1943, 1945, 1947; Wetherill 1975). For literature reviews of the early work in sequential analysis, see Jackson (1960) and Johnson (1961). SPRT sampling plans have been used in insect pest management to aid in monitoring insect populations or their damage since Stark (1952) developed the procedure for sampling the lodgepole needleminer, *Recurvaria milleri* Busck.

To develop the decision boundaries of the SPRT, the critical population levels for classification or decision-making (simple null and alternative hypotheses), probabilities (risk levels) for making Type I ( $\alpha$ ) and Type II ( $\beta$ ) errors, and underlying distribution of the variable or characteristic of interest are predetermined. Almost all SPRT insect sampling plans are based on the binomial, negative binomial, normal, and Poisson distributions.

In this paper, we review agriculture and forest entomology applications of the SPRT.

---

<sup>1</sup>School of Natural Resources, The University of Michigan, Ann Arbor, MI 48109-1115.

<sup>2</sup>School of Renewable Resources, The University of Arizona, Tucson, AZ 84721.

Table 1. Sequential probability ratio sampling plans for monitoring forest insect populations or their damage.

Insect	No. of SPRTs	$\alpha$ , $\beta$ values <sup>a</sup>
<b>Binomial Distribution</b>		
Larch sawfly (Ives 1954)	2	d
Larch sawfly (Ives and Prentice 1958)	2	d
Hemlock sawfly (Hard 1971)	1	a,d,e
Pine leaf adelgid (Dimond 1974)	2	d
Nantucket pine tip moth (Waters 1974)	1	a,d
<b>Negative Binomial Distribution</b>		
Spruce budworm (Morris 1954)	2	d
Spruce budworm (Waters 1955)	2	d
Forest tent caterpillar (Connola et al. 1957)	1	c
Red pine sawfly (Connola et al. 1959)	1	d
Spruce budworm (Cole 1960)	2	d
Spruce beetle (Knight 1960a)	1,2	d
Mountain pine beetle (Knight 1960b)	2	d,e
Cone and seed insects (Kozak 1964)	2	?
White grubs (Ives and Warren 1965)	1,2	d
Mountain pine beetle (Knight 1967)	2	c,d,e
Douglas-fir tussock moth (Mason 1969)	1	a,d
Spruce budworm (McKnight et al. 1970)	3	d,e,g
Roundheaded borers (Safranyik and Raske 1970)	2	d
Forest tent caterpillar (Shepherd and Brown 1971)	2	d
Jack pine sawfly (Tostowaryk and McLeod 1972)	2	d
Pine leaf adelgid (Dimond 1974)	2	d,f
Spruce budworm (Waters 1974)	2	d
<b>Normal Distribution</b>		
Lodgepole needleminer (Stark 1952)	2	b,d
Lodgepole needleminer (Stevens and Stark 1962)	3	d
Pine leaf adelgid (Dimond 1974)	2	d
<b>Poisson Distribution</b>		
Winter moth (Reeks 1956)	2	d
Spruce budworm (Cole 1960)	1	d
Saddled prominent eggs (Grimble and Kasile 1974)	1	c

<sup>a</sup>a=0.05, 0.05; b=0.05, 0.10; c=0.10, 0.05; d=0.10, 0.10; e=0.20, 0.20; f=0.30, 0.30; e=0.40, 0.40.

### PEST MANAGEMENT APPLICATIONS

Numerous sampling plans based on the SPRT for classifying various insect populations have been published in the entomology literature (Fowler and O'Regan 1974, Pieters 1978). At the present time, at least 65 applied articles have been published in insect pest management, 25 in forest entomology and 40 in agriculture entomology.

We summarize the applications in forest and agriculture entomology in tables 1 and 2, respectively, to include (1) the insect(s) sampled, (2) whether one or two SPRTs were used to develop a two-decision procedure (e.g., control versus no control) or a three-decision procedure (e.g., light versus medium versus heavy infestations), respectively, and (3) the mathematical distribution and values of  $\alpha$  and  $\beta$  used to develop the SPRT(s). Of the 28 SPRT sampling plans found in the 25 references in forest entomology,

Table 2. Sequential probability ratio sampling plans for monitoring agricultural insect populations or their damage.

Insect	No. of SPRTs	$\alpha, \beta$ values <sup>a</sup>
<b>Binomial Distribution</b>		
Green peach aphid, sugar beets (Sylvestor and Cox 1961)	1	f
Corn earworm, corn (Wolfenbarger and Darroch 1965)	1	f
Cotton fleahopper, cotton (Sterling and Pieters 1973)	3	f
Cotton fleahopper, cotton (Pieters and Sterling 1974)	1	f
Cotton arthropods, cotton (Sterling and Pieters 1974)	1	?
Bollweevil and <i>Heliothis</i> spp., cotton (Pieters and Sterling 1975)	2	f
Cotton arthropods, cotton (Sterling and Pieters 1975)	1	?
Cotton arthropods, cotton (Sterling 1976)	1	?
Mexican bean beetle, soybeans (Bellinger and Dively 1978)	1	e
Mexican bean beetle, soybeans (Bellinger et al. 1981)	1	b
<b>Negative Binomial Distribution</b>		
Green peach aphid, sugar beets (Sylvestor and Cox 1961)	1	f
Antestia, coffee trees (Rennison 1962)	1	c
Cabbage looper, cauliflower (Harcourt 1966a)	1	f
Imported cabbageworm, cauliflower (Harcourt 1966b)	2	f
Hylema brassicae (Bouché), cauliflower (Harcourt 1967)	1	f
Bollworm, cotton (Allen et al. 1972)	1	k
Boollworms, cotton (Ingram and Green 1972)	1	f
Lygus bugs, cotton (Sevacherian and Stern 1972)	1	f
Cabbage looper, cauliflower (Shepard 1973)	1	f
Alfalfa weevil, alfalfa (Stevens et al. 1976)	1	a,b,f
White grub, grain sorghum (Teetes and Sterling 1976)	1	?
<i>Culex tarsalis</i> Coquillett, rice (Mackey and Hoy 1978)	3	?
Hairy chinch bug, turfgrass and sod (Liu and McEwen 1979)	1	f
Cereal aphids, barley (Ba-Angood and Stewart 1980)	1	f
Corn rootworms, corn (Foster et al. 1982)	1	d
Tomato fruitworm, tomatoes (Nilakhe et al. 1982)	1	f,h,k
Japanese beetle, turfgrass and sod (Ng et al. 1983)	1	d,f
Alfalfa blotch leafminer, alfalfa (Harcourt 1983)	1	f
<b>Normal Distribution</b>		
Tomato pinworms, tomatoes (Wolfenbarger et al. 1975)	1	?
<b>Poisson Distribution</b>		
Asiatic rice borer, rice (Torii 1971)	2	?
<i>Aeneolamia varia saccharina</i> (Distant), sugar cane (Evans 1974)	1	f
Southern potato wireworm, potatoes (Onsager 1974)	1	i
<i>Nabis</i> spp., soybeans (Waddill et al. 1974)	2	f
<i>Geocoris</i> spp., soybeans (Waddill et al. 1974)	2	f
Wireworms, potatoes (Onsager et al. 1975)	2	g
Tomato pinworm, tomatoes (Wolfenbarger et al. 1975)	1	?
<i>Eurygaster integriceps</i> Put., winter wheat (Viktorov 1975)	1	f
Green cloverworm, soybeans (Hammond and Pedigo 1976)	1	f
Alfalfa weevil, alfalfa (Harcourt and Guppy 1976)	1	f
Velvetbean caterpillar, soybeans (Strayer et al. 1977)	2	f
Redbacked cutworm, peppermint (Danielson and Berry 1978)	1	j,l
Potato leafhopper, alfalfa (Luna et al. 1983)	1	d

<sup>a</sup>a=0.001, 0.001; b=0.01, 0.01; c=0.025, 0.025; d=0.05, 0.05; e=0.05, 0.01; f=0.10, 0.10; g=0.125, 0.125; h=0.15, 0.15; i=0.20, 0.05; j=0.20, 0.10; k=0.20, 0.20; l=0.40, 0.10.

5, 17, 3, and 3 were based on the binomial, negative binomial, normal, and Poisson distributions, respectively. Of the 42 SPRT sampling plans found in the 40 references in agriculture entomology, 10, 18, 1, and 13 were based on the binomial, negative binomial, normal, and Poisson distributions, respectively.

The proliferation of SPRT applications in the literature in recent years indicates widespread consideration of sequential sampling plans in insect pest management. In developing such plans, the sampler should be aware of the approximate nature of the decision process of Wald's SPRT due to overshooting errors and how such errors affect Wald's operating characteristic (OC) and average sample number (ASN) equations (Wald 1947). The errors caused by modifying the decision process, and the simultaneous operation of two or more SPRTs in a composite sampling plan should also be considered. For detailed discussions of these errors and how the sampler might adjust for them, see Fowler (1978, 1983, 1985).

#### ACKNOWLEDGMENTS

Our thanks to Dr. Frederick B. Knight, The University of Maine, Dr. Gary A. Simmons, Michigan State University, Dr. William E. Waters, University of California, and Dr. John A. Witter, The University of Michigan for critical review. This work was supported in part by the School of Natural Resources, The University of Michigan and the McIntire-Stennis Cooperative Forestry Research Program.

#### LITERATURE CITED

- Allen, J., D. Gonzalez, and D. V. Gokhale. 1972. Sequential sampling plans for the bollworm, *Heliothis zea*. Environ. Entomol. 1:771-780.
- Ba-Angood, S. A., and R. K. Stewart. 1980. Sequential sampling for cereal aphids on barley in southwestern Quebec. J. Econ. Entomol. 73:679-681.
- Bellinger, R. G., and G. P. Dively. 1978. Development of sequential sampling plans for insect defoliation on soybeans. J. New York Entomol. Soc. 86:278-279.
- Bellinger, R. G., G. P. Dively, II, and L. W. Douglass. 1981. Spatial distribution and sequential sampling of Mexican bean beetle defoliation on soybeans. Environ. Entomol. 10:835-841.
- Cole, W. E. 1960. Sequential sampling in spruce budworm control projects. For. Sci. 6:51-59.
- Connola, D. P., W. E. Waters, and W. E. Smith. 1957. The development and application of a sequential sampling plan for forest tent caterpillar in New York. New York State Mus. Bull. 366. 22 pp.
- Connola, D. P., W. E. Waters, and E. R. Nason. 1959. A sequential sampling plan for red-pine sawfly, *Neodiprion nanulus* Schedl. J. Econ. Entomol. 52:600-602.
- Danielson, S. D., and R. E. Berry. 1978. Redbacked cutworm: sequential sampling plans in peppermint. J. Econ. Entomol. 71:323-328.
- Dimond, J. B. 1974. Sequential surveys for the pine leaf chermid, *Pineus pinifoliae*. Univ. Maine. Life Sci. Agric. Exp. Sta. Tech. Bull. 68. 15 pp.
- Evans, D. E. 1974. Sequential sampling of adult sugar-cane froghoppers (*Aeneolamia varia saccharina* (Dist.)). Trop. Agric. 51:57-62.
- Foster, R. E., J. J. Tollefson, and K. L. Steffey. 1982. Sequential sampling plans for adult corn rootworms (Coleoptera: Chrysomelidae). J. Econ. Entomol. 75:791-793.
- Fowler, G. W. 1978. Errors in sampling plans based on Wald's sequential probability ratio test. Proc. 1977 Midwest Forest Mensurationists Meeting. USDA For. Serv. Gen. Tech. Rept. NC-46. 13 pp.
- \_\_\_\_\_. 1983. Accuracy of sequential sampling plans based on Wald's sequential probability ratio test. Canadian J. For. Res. 13:1197-1203.
- \_\_\_\_\_. 1985. The use of Wald's Sequential Probability Ratio Test to develop composite three-decision sampling plans. Canadian J. For. Res. 15:326-330.

- Fowler, G. W., and W. G. O'Regan. 1974. One-sided truncated sequential t-test: Application to natural resource sampling. USDA For. Serv. Res. Paper PSW-100. 17 pp.
- Ghosh, B. K. 1970. Sequential tests of statistical hypothesis. Addison-Wesley, Reading, Mass. 454 pp.
- Grimble, D. G., and J. D. Kasile. 1974. A sequential sampling plan for saddled prominent eggs. State Univ. New York, Coll. Environ. Sci. and For., Appl. For. Res. Inst. Rep. 15. 15 pp.
- Hammond, R. B., and L. P. Pedigo. 1976. Sequential sampling plans for green cloverworm in Iowa soybeans. J. Econ. Entomol. 69:181-185.
- Harcourt, D. G. 1966a. Sequential sampling for the imported cabbageworm, *Pieris rapae* (L.). Canadian Entomol. 98: 741-746.
- . 1966b. Sequential sampling for use in control of the cabbage looper on cauliflower. J. Econ. Entomol. 59:1190-1192.
- . 1967. Spatial arrangement of the eggs of *Hylema brassicae* (Bouché), and a sequential sampling plan for use in control of the species. Canadian J. Plant Sci. 47:461-467.
- . 1983. A sequential decision plan for management of the alfalfa blotch leafminer, *Agromyza frontella* (Diptera: Agromyziidae). Canadian Entomol. 115:1513-1518.
- Harcourt, D. G., and J. C. Guppy. 1976. A sequential decision plan for management of the alfalfa weevil, *Hypera postica* (Coleoptera: Curculionidae). Canadian Entomol. 108:551-555.
- Hard, J. S. 1971. Sequential sampling of hemlock sawfly eggs in southeast Alaska. USDA For. Serv. Res. Note PNW-142. 9 pp.
- Ingram, W. R., and S. M. Green. 1972. Sequential sampling for bollworms on raingrown cotton in Botswana. Cotton Grow. Rev. 49:265-275.
- Ives, W. G. H. 1954. Sequential sampling of insect populations. For. Chron. 30:287-291.
- Ives, W. G. H., and R. M. Prentice. 1958. A sequential sampling technique for surveys of the larch sawfly. Canadian Entomol. 90:331-338.
- Ives, W. G. H., and G. W. Warren. 1965. Sequential sampling for white grubs. Canadian Entomol. 97:596-604.
- Jackson, J. E. 1960. Bibliography on sequential analysis. J. Amer. Statist. Assoc. 55:561-580.
- Johnson, N. L. 1961. Sequential analysis: a survey. J. R. Statis. Soc. A 124:372-411.
- Knight, F. B. 1960a. Sequential sampling of Engelmann spruce beetle infestations in standing trees. USDA For. Serv. Res. Note RM-47. 4 pp.
- . 1960b. Sequential sampling of Black Hills beetle populations. U.S. For. Serv. Res. Note RM-48. 8 pp.
- . 1967. Evaluation of forest insect infestations. Ann. Rev. Entomol. 12:207-228.
- Kozak, A. 1964. Sequential sampling for improving cone collection and studying damage by cone and seed insects in Douglas fir. For. Chron. 40:210-218.
- Liu, H. J., and F. L. McEwen. 1979. The use of temperature accumulations and sequential sampling in predicting damaging populations of *Blissus leucopterus hirtus*. Environ. Entomol. 8:512-515.
- Luna, J. M., S. J. Fleischer, and W. A. Allen. 1983. Development and validation of sequential sampling plans for potato leafhopper (Homoptera: Cicadellidae) in alfalfa. Environ. Entomol. 12:1690-1694.
- Mackey, B. E., and J. B. Hoy. 1978. *Culex tarsalis*: Sequential sampling as a means of estimating populations in California rice fields. J. Econ. Entomol. 71:329-334.
- Mason, R. R. 1969. Sequential sampling of Douglas-fir tussock moth populations. USDA For. Serv. Res. Note PNW-102. 11 pp.
- McKnight, M. E., J. F. Chansler, D. B. Cahill, and H. W. Flake, Jr. 1970. Sequential plan for western budworm egg mass surveys in the central and southern rocky mountains. USDA For. Serv. Res. Note RM-174. 8 pp.
- Morris, R. F. 1954. A sequential sampling technique for spruce budworm egg surveys. Canadian J. Zool. 32:302-313.
- Ng, Y.-S., J. R. Trout, and S. Ahmad. 1983. Sequential sampling plans for larval populations of the Japanese beetle (Coleoptera: Scarabaeidae) in turfgrass. J. Econ. Entomol. 76:251-253.
- Nilakhe, S. S., R. B. Chalfant, S. C. Phatak, and B. Mullinix. 1982. Tomato fruitworm: Development of sequential sampling and comparison with conventional sampling in tomatoes. J. Econ. Entomol. 75:416-421.

- Oakland, G. B. 1950. An application of sequential analysis to whitefish sampling. *Biometrics* 6:59-67.
- Onsager, J. A. 1974. A sequential sampling plan for classifying infestations of southern potato wireworm. *Amer. Potato J.* 51:313-317.
- Onsager, J. A., B. J. Landis, and L. Fox. 1975. Efficacy of fonofos band treatments and a sampling plan for estimating wireworm populations in potatoes. *J. Econ. Entomol.* 68:199-202.
- Pieters, E. P. 1978. Bibliography of sequential sampling plans for insects. *Bull. Entomol. Soc. Amer.* 24:372-374.
- Pieters, E. P., and W. L. Sterling. 1974. A sequential sampling plan for the cotton fleahopper, *Pseudatomoscelis seriatus*. *Environ. Entomol.* 3:102-106.
- . 1975. Sequential sampling cotton squares damaged by boll weevils or *Heliothis* spp. in the coastal bend of Texas. *J. Econ. Entomol.* 68:543-545.
- Reeks, W. A. 1956. Sequential sampling for larvae of the winter moth, *Operophtera brumata* (Linn.) (Lepidoptera: Geometridae). *Canadian Entomol.* 88:241-246.
- Rennison, B. D. 1962. A method of sampling *Antestiopsis* in arabica coffee in chemical control schemes. *E. African Agric. For. J.* 27:197-200.
- Safranyik, L., and A. G. Raske. 1970. Sequential sampling plan for larvae of *Monochamus* in lodgepole pine logs. *J. Econ. Entomol.* 63:1903-1906.
- Sevacherian, V., and V. M. Stern. 1972. Sequential sampling plans for lygus bugs in California cotton fields. *Environ. Entomol.* 1:704-710.
- Shepard, M. 1973. A sequential sampling plan for treatment decisions on the cabbage looper on cabbage. *Environ. Entomol.* 2:901-903.
- Shepherd, R. F., and C. E. Brown. 1971. Sequential egg-band sampling and probability methods of predicting defoliation by *Malacasoma distria* (Lasiocampidae: Lepidoptera). *Canadian Entomol.* 103:1371-1379.
- Stark, R. W. 1952. Sequential sampling of the lodgepole needle miner. *For. Chron.* 28:57-60.
- Sterling, W. L. 1976. Sequential decision plans for the management of cotton arthropods in south-east Queensland. *Australian J. Ecol.* 1:265-274.
- Sterling, W. L., and E. P. Pieters. 1973. Sequential sampling plan for the cotton fleahopper. *Texas Agric. Ext. Serv. D-911.* 8 pp.
- . 1974. A sequential sampling package for key cotton arthropods in Texas. *Texas Agric. Exp. Sta. Tech. Rep.* 74-32. 28 pp.
- . 1975. Sequential sampling for key arthropods of cotton. *Texas Agric. Exp. Sta. Tech. Rep.* 75-24. 21 pp.
- Stevens, L. M., A. L. Steinhauer, M. J. Schlundt, R. F. Shroder, and J. U. McGuire. 1976. The biological meaning of sampling results and the development of a sequential sampling scheme to predict damaging levels of alfalfa weevil. *Univ. Maryland, Maryland Agric. Exp. Sta. MP-897.* 21 pp.
- Stevens, R. E., and R. W. Stark. 1962. Sequential sampling for the lodgepole needle miner, *Evagora milleri*. *J. Econ. Entomol.* 55:491-494.
- Strayer, J., M. Shepard, and S. G. Turnipseed. 1977. Sequential sampling for management decisions on the velvetbean caterpillar on soybeans. *J. Georgia Entomol. Soc.* 12:220-227.
- Sylvester, E. S., and E. L. Cox. 1961. Sequential plans for sampling aphids on sugar beets in Kern County, California. *J. Econ. Entomol.* 54:1080-1085.
- Teetes, G. L., and W. L. Sterling. 1976. A sequential sampling plan for a white grub in grain sorghum. *Southwest. Entomol.* 1:118-121.
- Torii, T. 1971. Quantitative occurrence prediction based on the sequential sampling test of the degrees of infestation by the rice stem borer. *Sci. Bull. Faculty of Agric., Kyushu Univ.* 25:103-112.
- Tostowaryk, W., and J. M. McLeod. 1972. Sequential sampling for egg clusters of the Swaine jackpine sawfly, *Neodiprion swainei* (Hymenoptera: Diprionidae). *Canadian Entomol.* 104:1343-1347.
- Viktorov, G. A. 1975. Method of sequential counting of the number of hibernating bugs of *Eurygaster integriceps*. *Soviet J. Ecol.* 6:278-280. (Translated from *Ékologiy* A6: 100-102, 1975).

- Waddill, V. H., B. M. Shepard, S. G. Turnipseed, and G. R. Carner. 1974. Sequential sampling plans for *Nabis* spp. and *Geocoris* spp. on soybeans. *Environ. Entomol.* 3:415-419.
- Wald, A. 1943. Sequential analysis of statistical data: Theory. Columbia Univ. Stat. Res. Grp. Rept. 75 and Off. Sci. Res. Develop. Rep. 1998.
- . 1945. Sequential tests of statistical hypotheses. *Ann. Math. Statist.* 16:117-186.
- . 1947. Sequential analysis. John Wiley & Sons, Inc., New York.
- Waters, W. E. 1955. Sequential sampling in forest insect surveys. *For. Sci.* 1:68-79.
- . 1974. Sequential sampling applied to forest insect surveys. IUFRO/SAF/SUNY Symposium on Monitoring Forest Environment Through Successive Sampling, June 24-26, 1974, Syracuse, NY. 22 pp.
- Wetherill, G. B. 1975. Sequential methods in statistics. John Wiley & Sons, Inc., New York. 218 pp.
- Wolfenbarger, D. A., and J. G. Darroch. 1965. A sequential sampling plan for determining the status of corn earworm control in sweet corn. *J. Econ. Entomol.* 58:651-654.
- Wolfenbarger, D. O., J. A. Cornell, S. D. Walker, and D. A. Wolfenbarger. 1975. Control and sequential sampling for damage by the tomato pinworm. *J. Econ. Entomol.* 68:458-460.



### INFORMATION FOR AUTHORS

Papers dealing with any aspect of entomology will be considered for publication in *The Great Lakes Entomologist*. Appropriate subjects are those of interest to amateur and professional entomologists in North Central States and Canada, as well as general papers and revisions directed to a larger audience while retaining an interest to readers in our geographical area.

Manuscripts must be typed, double-spaced, with wide margins on white 8-1/2 × 11" or equivalent size paper, and submitted in *duplicate*. Footnotes, legends, and captions of illustrations should be typed on separate sheets of paper. Titles should be concise, identifying the order and family discussed. The author of each species mentioned must be given fully at least once in the text. A common name for each species or group should be given at least once when such a name exists. The format of references should follow that used in recent issues. Photographs should be glossy. Drawings, charts, graphs, and maps must be scaled to permit proper reduction without loss of detail. Contributors should follow recommendations of the *Council of Biology Editors Style Manual*, 5th ed.

Papers published in *The Great Lakes Entomologist* are subject to a page charge of \$30.00 per published page. Members of the Society, who are authors without funds from grants, institutions, or industry and who are unable to pay costs from personal funds, may apply to the Society for financial assistance.

Authors will receive page proofs, together with an order blank for separates.

All manuscripts for *The Great Lakes Entomologist* should be sent to the Editor, Dr. D. C. L. Gosling, 69063 Wallowa Road, White Pigeon, Michigan 49099, USA. Other correspondence should be directed to the Executive Secretary (see inside front cover).