The Great Lakes Entomologist

Volume 22 Number 1 - Spring 1989 *Number 1 - Spring 1989*

Article 6

April 1989

An Annotated List of the Orthoptera of Beaver Island, Lake Michigan

R. G. Bland Central Michigan University

Follow this and additional works at: https://scholar.valpo.edu/tgle

Part of the Entomology Commons

Recommended Citation

Bland, R. G. 1989. "An Annotated List of the Orthoptera of Beaver Island, Lake Michigan," *The Great Lakes Entomologist*, vol 22 (1) DOI: https://doi.org/10.22543/0090-0222.1663 Available at: https://scholar.valpo.edu/tgle/vol22/iss1/6

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu. 1989

THE GREAT LAKES ENTOMOLOGIST

AN ANNOTATED LIST OF THE ORTHOPTERA OF BEAVER ISLAND, LAKE MICHIGAN

R. G. Bland¹

ABSTRACT

Thirty-six species of Orthoptera were collected from Beaver Island in northern Lake Michigan. Species distribution was Tetrigidae, 4; Acrididae, 16; Tettigoniidae, 8; Gryllacrididae, 1; and Gryllidae, 7.

Beaver Island (Charlevoix County, Michigan) is located approximately 52 km northwest of Charlevoix in northern Lake Michigan. It is 21 km long and 11 km wide with an area of 135 square km. The relatively undisturbed habitats include sand beaches with dune vegetation, upland fields, meadows, cedar swamps, bogs, and mixed deciduous and coniferous forests.

Little is known about orthopterans on islands of the Great Lakes. Pettit & McDaniel (1918) and Cantrall (1968) listed 15 species from Isle Royale but only 3 species from Beaver Island. Several references have included Michigan species as part of broader geographical studies (Blatchley 1920, Otte 1981 & 1984, Vickery & Kevan 1986) but do not refer to Beaver Island. The paucity of recorded species on the island was likely due to minimal collecting. Thus, the objective of this study was to determine which orthopteran species occurred on Beaver Island.

Collecting was done from 20 July to 7 August and 9 to 11 September 1987, and 4 to 6 June 1988. A few specimens collected by students in the 1960's and by the author in 1975 and 1980 were also recorded. A variety of habitats were sampled by sweeping and included dune vegetation, gravel pits, dry upland fields, agricultural fields, mowed grass, damp meadows, shorelines of lakes, shrubby edges of bogs, and mixed deciduous and coniferous woods. Pitfall traps containing molasses or antifreeze were placed in the above habitats and molasses-oatmeal trails were occasionally used. Katydids stridulating at night were stalked and captured.

Thirty-six species of Orthoptera were collected out of approximately 62 species listed from nearby counties of the Upper and Lower Peninsula of Michigan (Cantrall 1968, Vickery & Kevan 1986). An asterisk next to a species indicates a new Charlevoix County record.

Family TETRIGIDAE

Subfamily TETRIGINAE

Tetrix arenosa angusta (Hancock). 5 June-10 Sept. This commonly collected species occurred in the greatest diversity of habitats for tetrigids and was the most variable in

¹Biology Department, Central Michigan University, Mt. Pleasant, MI 48859

40

structure and colored markings, a characteristic common in this family (Rehn & Grant 1961).

Tetrix ornata ornata (Say). 4 June-10 Sept.

Tetrix subulata (L.). 5 June-15 Sept. This species was as abundant as T. arenosa angusta.

Subfamily BATRACHIDEINAE

Tettigidea lateralis (Say). 5 June-11 Sept. Nearly 85% of the specimens collected at Green's Lake in June were infested with larvae of an undescribed *Leptus* sp. (Erythraeidae), a bright pinkish-red mite. The infestation occurred in a damp to wet habitat dominated by two species of Cyperaceae (sedges): *Eleocharis* sp. (spike rush) and clumps of *Scirpus* sp. (bulrush). Iris (*Iris* sp.) were also present.

Tetrigids were encountered most frequently along the damp, flat shorelines of pools (e.g., in gravel pits, swamps), marshes, and shallow lakes. Shoreline vegetation was primarily short sedges and grasses, mosses, and organic debris. Tetrigids also frequented mixtures of damp, short, grass and broad-leaf vegetation growing on new logging trails at forest openings.

Family ACRIDIDAE

Subfamily GOMPHOCERINAE

Chloealtis conspersa (Harris). 20 July-6 August. Specimens were collected primarily in pitfall traps under trees along a field edge or in rank vegetation of fields.

Chorthippus curtipennis curtipennis (Harris). 12 July-11 Sept. This species was a very common acridid in lowland fields.

Pseudopomala brachyptera (Scudder). 20 July-2 August.

Subfamily OEDIPODINAE

Arphia pseudonietana pseudonietana (Thomas). 10 Sept. Groups of individuals occurred on extremely dry upland habitats consisting of sparse grass and broad-leaf herbs or grass and reindeer lichen.

Chortophaga viridifasciata (DeGeer), 4 June–12 August. This was the only adult acridid observed in early June except for a few freshly-molted Melanoplus borealis borealis (Fieber).

Camnula pellucida (Scudder). 20 July-11 Sept. This acridid was the most abundant species; it occurred in all upland, moist lowland, and agricultural fields.

Dissosteira carolina (L.). 21 July-7 August. Spharagemon collare (Scudder). 7 August-10 Sept. The few individuals observed occurred in a small corn field and a recently abandoned, weedy agricultural field.

Trimerotropis huroniana E. M. Walker. 20 July-11 Sept. Individuals frequented the sparsely vegetated sand dunes of Lake Michigan shorelines.

Subfamily MELANOPLINAE

Booneacris glacialis canadensis (E. M. Walker). 3 August-11 Sept. Individuals were captured on leatherleaf along the birch and maple edge of the nearly dry, remnant bog of Green's Lake.

Melanoplus bivittatus (Say). 22 July-5 August.

Melanoplus borealis borealis (Fieber). 4 June-6 August. Individuals were netted or

https://scholar.valpo.edu/tgle/vol22/iss1/6 DOI: 10.22543/0090-0222.1663

Bland: An Annotated List of the Orthoptera of Beaver Island, Lake Michig

1989

THE GREAT LAKES ENTOMOLOGIST

41

collected from pitfall traps in rank vegetation of fields or under trees at field edges. The first adults were collected 14 days before the earliest collecting record listed by Cantrall (1968).

Melanoplus confusus Scudder*. July 23.

Melanoplus femurrubrum femurrubrum (DeGeer). 27 July-10 Sept.

Melanoplus islandicus Blatchley. 30 July–11 Sept. Individuals were captured in a habitat of grass, broad-leaf ground cover, and moss at forest edges, as well as sunlit forest openings alongside a trail above Iron Ore Creek.

Melanoplus sanguinipes sanguinipes (F.). 20 July-10 Sept. This species reached a peak in late August and early September to become nearly as abundant as C. pellucida.

Cantrall (1968) listed approximately 14 species found in nearby mainland counties that might be expected to occur on Beaver Island but were not collected in my study. In particular, the yellow wings and crepitation of *A. sulphurea* (F.) would have made this species easy to locate in June if present, and the same for *S. bolli* Scudder in sandy fields, roadsides, or forest edges in August and September. *Trimerotropis verruculata* (Wm. Kirby) would have produced loud crepitations around gravel pits near wooded areas (Vickery & Kevan 1986). *Orphulella speciosa* (Scudder), *M. dawsoni* (Scudder), and *M. keeleri luridus* (Dodge) were missing species common on the sandy, grassy wasteland of the surrounding mainland, and *M. fasciatus* was not found in sunlit, woodland openings of *Vaccinium* or other low shrubs as expected. *Encoptolophus sordidus* (Burmeister) is a common September mainland species also not observed.

Family TETTIGONIIDAE

Subfamily PHANEROPTERINAE

Scudderia curvicauda (DeGeer)*. 20 July–10 Sept. Scudderia furcata furcata Brunner. 31 July–25 Sept. Scudderia pistillata Brunner. 20 July–10 Sept. Scudderia spp. were collected primarily from shrubs and broad-leaf herbs of upland and

lowland grassy fields.

Subfamily COPIPHORINAE

Neoconocephalus ensiger (Harris)*. 29 July-10 Sept.

Subfamily CONOCEPHALINAE

Conocephalus fasciatus (DeGeer). 22 July-10 Sept. This species ranged from dry, upland fields to damp, shoreline vegetation.

Conocephalus saltans (Scudder). 21 July-10 Sept. Individuals occurred in dry, upland fields.

Orchelimum gladiator Bruner. 20 July-30 Sept. This species was most common in damp meadows of tall sedges and grasses.

Subfamily DECTICINAE

Atlanticus monticola Davis^{*}. 20 July-22 Sept. This species was collected from low branches of small hemlock and oak trees in open areas, as well as pitfall traps placed among bracken ferns near maple woods, under a maple tree, and in a weedy corn field. The other Michigan species, *A. testaceous* (Scudder), was not found.

42

THE GREAT LAKES ENTOMOLOGIST

Vol. 22, No. 1

Family GRYLLACRIDIDAE

Subfamily RHAPHIDOPHORINAE

Ceuthophilus maculatus (Harris)*. 17 July-7 August. Several individuals were collected at night from inside an old stump. Otherwise, the same pit traps that captured *A. monitcola*, and another next to a fallen log, yielded most of the specimens. Four additional species, *C. brevipes* Scudder in particular, may occur on the island based on their ranges (Cantrall 1968).

Family GRYLLIDAE

Subfamily GRYLLINAE

Gryllus pennsylvanicus Burmeister. 22 July-11 Sept. Gryllus veletis (R. D. Alexander & Bigelow). 5 June-2 July.

Subfamily NEMOBIINAE

Allonemobius (= Nemobius) allardi (R. D. Alexander & E. S. Thomas). 21 July-30 Sept. The first adult was collected 7 days before the earliest date listed by Cantrall (1968). Allonemobius (= Nemobius) fasciatus (DeGeer). 31 July-10 Sept. Allonemobius (= Nemobius) griseus griseus (E. M. Walker). 2 August-10 Sept. Eunemobius (= Nemobius) carolinus carolinus (Scudder). 11 Sept.

Subfamily OECANTHINAE

Oecanthus quadripunctatus Beutenmüller. 5 August-10 Sept.

The six species of Gryllinae and Nemobilinae represent all species naturally occurring in northern Michigan. They were collected from pitfall traps in a variety of upland and lowland habitats. O. quadripunctatus was common on low shrubs and broad-leaf herbs of fields and roadsides. O. nigricornis F. Walker was not collected but its habitats (Vickery & Kevan 1986) occur on the island and most likely it is present.

ACKNOWLEDGMENTS

D. Profant, Biology Department, Central Michigan University (C.M.U.), assisted in collecting insects. V. Vickery, Lyman Entomological Museum of McGill University, Quebec, confirmed the identification of several species, and W. Welbourn, Acarology Laboratory, Ohio State University, identified the mite species. This study was funded by a C.M.U. faculty research grant and facilities were made available by the C.M.U. Biological Station, Beaver Island.

LITERATURE CITED

Blatchley, W. S. 1920. Orthoptera of Northeastern America. Nature Publ. Co., Indianapolis. 784 pp. Cantrall, I. J. 1968. An annotated list of the Dermaptera, Dictyoptera, Phasmatoptera, and Orthoptera of Michigan. Mich. Entomol. 1:299-346.

Otte, D. 1981. The North American Grasshoppers. Vol. 1. Acrididae, Gomphocerinae and Acridinae. Harvard Univ. Press, Cambridge, MA. 275 pp.

https://scholar.valpo.edu/tgle/vol22/iss1/6 DOI: 10.22543/0090-0222.1663

Bland: An Annotated List of the Orthoptera of Beaver Island, Lake Michig

1989

THE GREAT LAKES ENTOMOLOGIST

43

_____. 1984. The North American Grasshoppers. Vol. II. Oedipodinae. Harvard Univ. Press, Cambridge, MA. 366 pp.

Pettit, R. H. and E. McDaniel. 1918. Key to Orthoptera of Michigan with annotations. Mich. Agr. Coll. Spec. Bull. 83. 48 pp.

Rehn, J. A. G. and H. J. Grant, Jr. 1961. A monograph of the Orthoptera of North America (North of Mexico). Vol. I. Mon. Acad. Nat. Sci. Phila. no. 12. 257 pp. + 8 pl.

Vickery, V. R. and D.K. McE. Kevan. 1986. The insects and arachnids of Canada. Part 14. The grasshoppers, crickets and related insects of Canada and adjacent regions. Ulonata: Dermaptera, Cheleutoptera, Notoptera, Dictyoptera, Grylloptera and Orthoptera. Agric. Canada Publ. 1777 (1985).