

December 1991

Orientation of *Hylobius Pales* and *Pachylobius Picivorus* (Coleoptera: Curculionidae) to Visual Cues

D.W. A. Hunt
University of Wisconsin

K. F. Raffa
University of Wisconsin

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

Hunt, D.W. A. and Raffa, K. F. 1991. "Orientation of *Hylobius Pales* and *Pachylobius Picivorus* (Coleoptera: Curculionidae) to Visual Cues," *The Great Lakes Entomologist*, vol 24 (4)
Available at: <https://scholar.valpo.edu/tgle/vol24/iss4/3>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

ORIENTATION OF *HYLOBIUS PALES* AND *PACHYLOBIUS PICIVORUS* (COLEOPTERA: CURCULIONIDAE) TO VISUAL CUES

D. W. A. Hunt,^{1,2} and K. F. Raffa¹

ABSTRACT

Pitfall traps with above-ground silhouettes of various colors and diameters were used in field tests to evaluate the role of vision in host orientation by adult pales weevils, *Hylobius pales*, and pitch-eating weevils, *Pachylobius picivorus*. White traps (11 cm outer diameter) baited with ethanol and turpentine caught significantly more weevils than similarly baited black or green traps (11 cm outer diameter). Trap diameter (range of 6–22 cm outer diameter) did not affect trap catch. Pitfall traps can be used to monitor root weevil populations in young pine plantations and Christmas tree farms, where they are major pests. These results demonstrate that visual and chemical cues can be integrated to improve trap efficiency.

The pales weevil, *Hylobius pales* (Herbst), and the pitch-eating weevil, *Pachylobius picivorus* (Germar), are important pests of a variety of pine species in eastern North America. The adults of both species feed on the inner bark of young pine stems, and can cause widespread mortality in recently planted pine plantations (Nord et al. 1982). Adult feeding on stems and branches can also disfigure young trees (Drooz 1985). Along with the pine root collar weevil, *Hylobius radialis* Buchanan, these root weevils comprise the most important pest group affecting Christmas trees in Wisconsin. Infestations are difficult to detect, so growers rely on routine applications of persistent insecticides, primarily Lindane.

Recent research has focused on the development of a simple trap-based monitoring technique for forecasting weevil damage (Hunt and Raffa 1989, Rieske and Raffa 1990). The objective is to replace calendar applications with threshold-triggered treatments to reduce insecticide use, and allow for conversion to less persistent materials. Monitoring efficiency is increased through the use of cues that weevils use to orient to host trees. A variety of host odors appear to be involved in host location by adult weevils. Host monoterpenes are attractive to *H. pales* (Thomas and Hertel 1969, 1979), and *P. picivorus* (Fatzinger 1985). The combination of ethanol and terpenes is also attractive to both species, and factors such as ratios of the components and season can greatly affect attraction (Fatzinger et al. 1987, Raffa and Hunt 1988, Hunt and Raffa 1989, Rieske and Raffa 1991). In contrast, relatively little is known about the possible role of visual cues in host orientation. It was our goal to determine whether visual cues influence orientation by *H. pales* and *P. picivorus* to attractant-baited traps.

¹Department of Entomology, University of Wisconsin, Madison, WI 53706.

²Current address: Agriculture Canada, Research Station, Harrow, Ontario, N0R 1G0, Canada.

MATERIALS AND METHODS

The traps used were a modified version of the pitfall trap described by Tilles et al. (1986a, b) and modified by Hunt and Raffa (1989). Sections of PVC drainpipe, 76 cm long, were drilled with eight 6 mm diameter holes spaced equally around the circumference of the pipe, 31 cm from one end. The pipes were inserted into the ground to a depth of 31 cm so that the holes in the pipe were at ground level, and the remaining 46 cm provided a silhouette. The pipes were capped at both ends with removable plastic lids, with two 0.3 cm diameter holes in the lower lid to allow water drainage. Liquid Teflon was applied to the inner surfaces of the traps to prevent the escape of weevils. A wire passing through two 2 mm holes in the walls of the trap at ground level served as a support for baits.

To examine the effect of trap color, the above ground portion of 11 cm outer diameter traps was painted black, white or green using a flat spray enamel (New York Bronze Powder Co. Inc., Elizabeth, NJ). To examine the effect of diameter, black PVC of 22 or 6 cm outer diameter were compared. These sizes were used so that the weevils could choose between traps of greater and lesser diameter than the trees present in the plantation. The traps were placed in a 4 yr old Scotch pine Christmas tree plantation in Waushara County, WI. The plantation was known to be infested with *H. pales* and *P. picivorus*. The trees had been planted at 1.7 m between trees and rows. The traps were placed between the rows, every third row, midway between every third tree. This resulted in approximately a 5.1 x 5.1 m spacing of groups of traps. The traps were arranged in 20 pairs for the diameter experiment and at the points of 20 equilateral triangles for the color experiment, with 0.6 m between traps in a group. The position of each color or diameter of trap was assigned randomly within each triplet or pair, respectively.

The baits consisted of vials of turpentine (Sunnyside Corp., Wheeling, IL) and 95% ethanol. The turpentine and 95% ethanol were each released separately from open 12 x 35 mm glass vials, such that the release rates were approximately 40 mg and 400 mg/24 h, respectively, at 22° C. Vials within each type of trap exhibited different rates of evaporation, probably due to temperature differences within the trap which were characteristic of each color and diameter of PVC tube. In order to standardize the release rates a variable number of vials was used for each color and diameter of trap. The turpentine consisted mostly of monoterpenes, of which the relative proportions were: alpha-pinene (52.45%), beta-pinene (41.35%), beta-phellandrene (2.00%), limonene (1.05%), camphene (0.85%), myrcene (0.65%), and unknown (0.65%), as determined by gas liquid chromatography using the method of Raffa and Steffeck (1988). Weevils were collected from the traps and the baits replenished at approximately weekly intervals from May to Sept., 1987. All weevils were examined in the laboratory for species identity using the keys developed by Kissinger (1964) and Warner (1966), and sex was determined by the methods of Wilson et al. (1966).

Numbers of weevils of each species captured in pitfall traps were analyzed using analysis of variance and Duncan's multiple range test. Statistical Analysis System programs were used for all analyses (SAS Institute 1982).

RESULTS AND DISCUSSION

Adult *H. pales* and *P. picivorus* of both sexes were captured in significantly larger numbers in baited pitfall traps that were painted white than in similarly baited pitfall traps painted black or green (Table 1). This seemed surprising because black or green traps more closely approximate the color of the host trunk and foliage, respectively. However, *H. pales* and *P. picivorus* breed primarily in the stumps and roots of pines that have recently been harvested or were killed by bark beetles, fire, or some other agent (Nord et al. 1982), so these insects may not be very responsive to

Table 1. — Mean number (\pm SE) of *Hylobius pales* and *Pachylobius picivorus* adults caught per trap from 19 May to 7 Sept. 1987, in 11 cm diameter pitfall traps of various colors in Wisconsin. N = 20 traps for each color^a.

| Trap color | <i>H. pales</i> | | <i>P. picivorus</i> | |
|----------------|-----------------|----------------|---------------------|----------------|
| | Females | Males | Females | Males |
| White | 17.3 \pm 3.2a | 1.8 \pm 0.7a | 15.3 \pm 3.9a | 9.8 \pm 2.3a |
| Black | 2.5 \pm 0.8b | 0.5 \pm 0.3b | 2.8 \pm 1.1b | 1.8 \pm 0.8b |
| Green | 0.5 \pm 0.03b | 0 b | 0 b | 0 b |
| Total captured | 406 | 46 | 362 | 232 |

^aMeans within columns followed by the same letter are not significantly different ($P = 0.05$; Duncan's multiple range test [SAS Institute 1982]).

Table 2. — Mean number (\pm SE) of *Hylobius'pales* and *Pachylobius picivorus* adults caught per trap from 19 May to 7 Sept. 1987, in black pitfall traps of different diameters (large = 22 cm; small = 6 cm) in Wisconsin. N = 20 traps for each diameter^a.

| Trap diameter | <i>H. pales</i> | | <i>P. picivorus</i> | |
|----------------|-----------------|----------------|---------------------|----------------|
| | Females | Males | Females | Males |
| Large | 3.2 \pm 1.0a | 0.7 \pm 0.4a | 3.1 \pm 0.9a | 1.7 \pm 0.7a |
| Small | 2.9 \pm 0.8a | 0.8 \pm 0.4a | 3.0 \pm 0.8a | 1.5 \pm 0.5a |
| Total captured | 122 | 30 | 122 | 64 |

^aMeans within columns followed by the same letter are not significantly different ($P = 0.05$; Duncan's multiple range test [SAS Institute 1982]).

host color. Also, both of these species feed nocturnally, so white traps may simply provide greater contrast at night.

Our data on the effects of color on trap catch contradicts the results of Fatzinger (1985), who reported that in Florida more *H. pales* and *P. picivorus* were caught in black flight-barrier traps than in white traps. This discrepancy may be due to differences in the behavior of flying and walking weevils, or may reflect differences in Wisconsin and Florida populations.

Trap diameter did not affect catch for either *H. pales* or *P. picivorus* (Table 2). Orientation to an odor source may be a more effective method than diameter-related preference for locating stumps and roots, since a silhouette may be absent or of variable dimensions above suitable host material. For these weevils tree condition may determine host suitability more than age or size. Similarly, Nordlander et al. (1986) reported that *Hylobius abietis* oriented to underground odor sources in the absence of visual cues. The large proportion of female weevils caught in traps baited with ethanol combined with turpentine is in agreement with Hunt and Raffa (1989), and suggests that these compounds are used as chemical cues primarily by females.

Weevil orientation to baited traps of various colors in this study suggests that visual cues play a role in host orientation by root weevils. However, neither *P. picivorus* nor *H. radialis* are attracted to black pitfall traps in the absence of turpentine and ethanol baits (Hunt and Raffa 1989), suggesting that visual orientation can augment olfactory orientation but not replace it. Because *H. radialis* breed in the root collar area of standing pine trees (Schaffner and McIntyre 1944), as opposed to the stumps and roots used by *H. pales* and *P. picivorus*, visual cues such as color or diameter may be even more important in *H. radialis*.

Previous studies of host orientation in pine root weevils have used black pitfall traps (Raffa and Hunt 1988, Hunt and Raffa 1989). Our results suggest that more weevils would have been captured if these studies had used a more attractive or more visible color such as white. Since the number of certain species of root weevils

captured in pitfall traps appears to be correlated with tree damage in pine plantations (Nordlander 1987, Hunt and Raffa 1989) the use of white traps could increase trapping efficiency, and thereby provide for a more sensitive and accurate monitoring tool.

ACKNOWLEDGMENTS

This work was supported in part by USDA grants CRCR-1-2077 and FSTY-9-0210, the Wisconsin Department of Natural Resources, the University of Wisconsin-Madison College of Agricultural and Life Sciences, the University of Wisconsin Applied Research Program, the Christmas Tree Producers Associations of Wisconsin, Michigan, Minnesota, and Illinois, and a Natural Sciences and Engineering Research Council Postdoctoral Fellowship to the senior author. We thank Brian Strom for trap construction, Dr. David Hall of the Wisconsin Department of Natural Resources and Lorin Zastoupil of the Kirk Company for their valuable advice. We also thank the Kirk Company (Wautoma, WI) for providing the study site.

LITERATURE CITED

- Drooz, A. T. [ed.] 1985. Insects of eastern forests. USDA For. Serv. Misc. Pub. 1426. Washington, D.C.
- Fatzinger, C. W. 1985. Attraction of the black turpentine beetle (Coleoptera: Scolytidae) and other forest Coleoptera to turpentine-baited traps. *Environ. Entomol.* 14:768-775.
- Fatzinger, C. W., B. D. Siegfried, R. C. Wilkinson and J. L. Nation. 1987. Trans-verbenol, turpentine, and ethanol as trap baits for the black turpentine beetle, *Dendroctonus terebrans*, and other forest Coleoptera in North Florida. *J. Entomol. Sci.* 22:201-209.
- Hunt, D. W. A. and K. F. Raffa. 1989. Attraction of *Hylobius radicis* and *Pachylobius picivorus* (Coleoptera: Curculionidae) to ethanol and turpentine in pitfall traps. *Environ. Entomol.* 18:351-355.
- Kissinger, D. G. 1964. Curculionidae of America north of Mexico: A key to the genera. Taxonomic Publications, South Lancaster, Massachusetts.
- Nord, J. C., J. H. Ghent, H. A. Thomas and C. A. Doggett. 1982. Control of pales and pitch-eating weevils in the south. USDA For. Serv. For. Rep. SA-FR 21.
- Nordlander, G. 1987. A method for trapping *Hylobius abietis* (L.) with a standardized bait and its potential for forecasting seedling damage. *Scand. J. For. Res.* 2:199-213.
- Nordlander, G., H. H. Eidmann, U. Jacobsson, H. Nordenhem and K. Sjödin. 1986. Orientation of the pine weevil *Hylobius abietis* to underground sources of host volatiles. *Entomol. exp. appl.* 41:91-100.
- Raffa, K. F. and D. W. A. Hunt. 1988. Use of baited pitfall traps for monitoring pales weevil, *Hylobius pales* (Herbst) (Coleoptera: Curculionidae), adult populations. *Great Lakes Entomol.* 21:123-125.
- Raffa, K. F. and R. J. Steffeck. 1988. Computation of response factors for quantitative analysis of monoterpenes by gas-liquid chromatography. *J. Chem. Ecol.* 14:1385-1390.
- Rieske, L. K. and K. F. Raffa. 1990. Use of a monitoring system to evaluate pesticide efficacy and residual activity against two pine root weevils, *Hylobius pales* and *Pachylobius picivorus* (Coleoptera: Curculionidae), in Christmas tree farms. *Great Lakes Entomol.* 23:189-193.
- Rieske, L. K. and K. F. Raffa. 1991. Effect of varying ethanol and turpentine ratios on attraction of two pine root weevil species, *Hylobius pales* and *Pachylobius picivorus* (Coleoptera: Curculionidae). *Environ. Entomol.* 20:48-52.
- SAS Institute. 1982. SAS user's guide: statistics. SAS Institute, Cary, N.C.
- Schaffner, J. V., Jr. and H. I. McIntyre. 1944. The pine root collar weevil. *J. For.* 42:269-275.

- Thomas, H. A. and G. D. Hertel. 1969. Responses of the pales weevil to natural and synthetic host attractants. *J. Econ. Entomol.* 62:383-386.
- Thomas, H. A. and G. D. Hertel. 1979. Responses of pales weevils to natural and synthetic traps under field conditions. *J. Econ. Entomol.* 72:342-345.
- Tilles, D. A., K. Sjodin, G. Nordlander and H. H. Eidmann. 1986a. Synergism between ethanol and conifer host volatiles as attractants for the pine weevil, *Hylobius abietis* (L.) (Coleoptera: Curculionidae). *J. Econ. Entomol.* 79:970-973.
- Tilles, D. A., G. Nordlander, H. Nordheim, H. H. Eidmann, A. B. Wassgren and G. Bergstrom. 1986b. Increased release of host volatiles from feeding scars: A major cause of field aggregation in the pine weevil *Hylobius abietis* (Coleoptera: Curculionidae). *Environ. Entomol.* 15:1040-1054.
- Warner, R. E. 1966. A review of the *Hylobius* of North America, with a new species injurious to slash pine (Coleoptera: Curculionidae). *Coleop. Bull.* 20:65-81.
- Wilson, L. F., C. D. Waddell and I. Millers. 1966. A way to distinguish sex of adult *Hylobius* weevils in the field. *Can. Entomol.* 98:1118-1119.