

The Great Lakes Entomologist

Volume 25
Number 2 - Summer 1992 *Number 2 - Summer*
1992

Article 10

June 1992

The Sand-Dwelling Predatory Mayfly *Pseudiron Centralis* in Michigan (Ephemeroptera: Pseudironidae)

W. P. McCafferty
Purdue University

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

McCafferty, W. P. 1992. "The Sand-Dwelling Predatory Mayfly *Pseudiron Centralis* in Michigan (Ephemeroptera: Pseudironidae)," *The Great Lakes Entomologist*, vol 25 (2)
DOI: <https://doi.org/10.22543/0090-0222.1779>
Available at: <https://scholar.valpo.edu/tgle/vol25/iss2/10>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

THE SAND-DWELLING PREDATORY MAYFLY *PSEUDIRON CENTRALIS* IN MICHIGAN (EPHEMEROPTERA: PSEUDIRONIDAE)¹W. P. McCafferty²

An examination of samples of Ephemeroptera larvae taken from 19 sites on the Pere Marquette River, Mason County, Michigan in October, 1990 and May, 1991 by A. G. B. Primack (Indiana University, Bloomington) revealed the presence of *Pseudiron centralis* McDunnough. This species was taken at seven sites between State Highway M37 and Rainbow Rapids, east of Ludington. Voucher specimens are held at the Purdue Entomological Research Collection. Michigan has been surveyed extensively for riverine benthic macroinvertebrates in the past, and mayflies in particular are relatively well documented in Michigan because of their importance to fly fishing (e.g., Leonard and Leonard 1962), bioassessment, etc. Despite these facts, however, this distinctive and somewhat large-sized mayfly has not been previously reported for the state.

The new record is noteworthy because *Pseudiron centralis* is an unusual and fascinating species that may be endangered in parts of its range. Larvae are sand-dwelling midge hunters (Clifford and Soluk 1985, Soluk and Craig 1990) that move along the sand much like crabs (Clifford and Soluk 1984) but are also capable of extremely fast swimming (McCafferty and Provonsha 1986). It is the only recognized species of the Nearctic family Pseudironidae (Pescador 1985, McCafferty 1991a), and its distribution in the southeastern, central, and some northwestern United States and west-central Canada is typical of many unrelated psammophilous mayflies (McCafferty 1991b). Michigan represents the most northeastern record for the species. It is known from nearby states of Indiana, Illinois, and Wisconsin. In Indiana, it has recently been considered an endangered species by that state's Natural Resources Commission (Anonymous 1992).

Besides sand and sand/gravel habitats, a few larvae of *P. centralis* were taken on silt, in *Elodea* beds, on logs, or under cut banks on the Pere Marquette River. This may indicate a broader habitat preference for the species than previously thought, or, more likely, that the larvae are prone to drift and thus can be found in atypical habitats.

Pescador (1985) noted a distinct geographic gradation with regard to degree of body pigmentation found in populations of *P. centralis*, with individuals of northern populations being darker than those of southern populations. The specimens from the Pere Marquette River are typical of northern populations. The frons and vertex of the head of the Michigan specimens are dark brown and therefore, according to Pescador (1985), are not typical of other populations from central continental areas, but instead are typical of populations from Alberta, Saskatchewan, Utah, and Wyoming.

¹Published as Purdue Experiment Station Journal No. 13397.

²Department of Entomology, Purdue University, West Lafayette, IN 47907.

LITERATURE CITED

- Anonymous. 1992. Indiana Department of Natural Resources nonrule policy document. Nat. Res. Comm. Inform. Bull. No. 2. Ind. Register Vol. 15, No. 7:1312-1327.
- Leonard, J. W. and F. A. Leonard. 1962. Mayflies of Michigan trout streams. Cranbrook Inst. Sci., Bloomfield, Mich.
- McCafferty, W. P. and A. V. Provonsha. 1986. Comparative mouthpart morphology and evolution of the carnivorous Heptageniidae (Ephemeroptera). Aquatic Insects 8:83-89.
- McCafferty, W. P. 1991a. Toward a phylogenetic classification of the Ephemeroptera (Insecta): a commentary on systematics. Ann. Entomol. Soc. Am. 84:343-360.
- _____. 1991b. Comparison of Old and New World *Acanthametropus* (Ephemeroptera: Acanthametropodidae) and other psammophilous mayflies. Entomol. News 102:205-214.
- Pescador, M. L. 1985. Systematics of the Nearctic genus *Pseudiron* (Ephemeroptera: Heptageniidae: Pseudironinae). Fla. Entomol. 68:432-444.
- Soluk, D. A. and H. F. Clifford. 1984. Life history and abundance of the predaceous psammophilous mayfly *Pseudiron centralis* McDunnough (Ephemeroptera: Heptageniidae). Can. J. Zool. 62:1534-1539.
- _____. 1985. Microhabitat shifts and substrate selection by the psammophilous predator *Pseudiron centralis* McDunnough (Ephemeroptera: Heptageniidae). Can. J. Zool. 63:1539-1543.
- Soluk, D. A. and D. A. Craig. 1990. Digging with a vortex: flow manipulation facilitates prey capture by a predatory stream mayfly. Limnol. Oceanogr. 35:1201-1206.