

# The Great Lakes Entomologist

---

Volume 40

Number 3 & 4 - Fall/Winter 2007 *Number 3 & 4 -  
Fall/Winter 2007*

Article 7

---

October 2007

## Parasitism of *Urophora Affinis* (Diptera: Tephritidae) by *Aprostocetus* Sp. (Hymenoptera: Eulophidae) in Michigan

Jordan M. Marshall

*Michigan Technological University*

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

---

### Recommended Citation

Marshall, Jordan M. (2007) "Parasitism of *Urophora Affinis* (Diptera: Tephritidae) by *Aprostocetus* Sp. (Hymenoptera: Eulophidae) in Michigan," *The Great Lakes Entomologist*: Vol. 40 : No. 2 , Article 7.

Available at: <https://scholar.valpo.edu/tgle/vol40/iss2/7>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

**PARASITISM OF *UROPHORA AFFINIS* (DIPTERA: TEPHRITIDAE)  
BY *APROSTOCETUS* SP. (HYMENOPTERA: EULOPHIDAE)  
IN MICHIGAN**

Jordan M. Marshall<sup>1</sup>

*Urophora affinis* Frfld. and *U. quadrifasciata* (Meig.) (Diptera: Tephritidae) are Eurasian gallicolous fruit flies introduced to North America in 1972 as biological control agents for *Centaurea biebersteinii* DC (spotted knapweed, Asteraceae, = *C. maculosa* auct. non Lam.) (Harris 1980). Through natural dispersal and numerous introductions, both *Urophora* species have become distributed throughout the introduced range of *C. biebersteinii* (Foote et al. 1993, Lang et al. 1997, Mays and Kok 2003). Use as biological control agents for *C. biebersteinii* focused on the diversion of energy from seed production to the development of *Urophora* spp. larvae within galls (Harris 1980). In the introduced range of *U. affinis* and *U. quadrifasciata*, mortality of these species has occurred as a result of predation by bird and mammal species, as well as parasitism by *Pteromalus cardui* (Erdős) (Hymenoptera: Pteromalidae) (Story et al. 1995, Pearson 1999, Marshall et al. 2004).

*Aprostocetus* sp. (Hymenoptera: Eulophidae) was initially observed emerging from *C. biebersteinii* seed heads in rearing during February 2008. Typically the subfamily Tetrastichinae, containing the genus *Aprostocetus* Westwood, is endoparasitic of eggs, larvae, or pupae of Coleoptera, Diptera, Hymenoptera, and Lepidoptera, with a distinct association with gall inducing hosts (Noyes 2003, Yegorenkova et al. 2007). Species within *Aprostocetus*, approximately 670, are distributed globally (Graham 1987, LaSalle 1994, Yegorenkova et al. 2007).

*Centaurea biebersteinii* seed heads were collected from three sites in Livingston County, MI, on 18 February 2008. Sixty seed heads from each site were randomly selected and placed into 8 dram plastic shell vials for rearing. Vials were a third filled with wet sand, topped with a layer of dry sand, and were capped with cotton fabric. Each vial contained 2 *C. biebersteinii* seed heads (30 vials/site), stored at 80° C at 45 percent humidity, and were checked every 2 days until emergence began. Following initial emergence, vials were checked daily and *Aprostocetus* sp. adults were removed and sexed. Adult *Urophora* species were identified using Foote et al. (1993). Chi-square tests were used to test for the independence of *U. affinis* from the occurrence of *Aprostocetus* sp. A *t*-test was used to test for differences in *U. affinis* emergence in the presence of *Aprostocetus* sp.

*Aprostocetus* sp. first emerged after 14 days. Seed heads were left in vials for an additional 10 weeks, which resulted in 34.4 percent of vials producing a total of 373 adult *Aprostocetus* sp. All emergence of *Aprostocetus* sp. occurred during days 14-17. In vials that produced adult *Aprostocetus* sp., 12.03 ± 9.04 individuals emerged per vial with a sex ratio of 0.36 males/females. A total of 36 adult *U. affinis* emerged from 26.7 percent of vials with a sex ratio of 0.89 males/females. In vials that produced adult *U. affinis*, 1.50 ± 0.83 individuals emerged per vial. The presence of *U. affinis* was independent of the presence of *Aprostocetus* sp. individuals ( $\chi^2 = 0.404$ , *df* = 1). There was no significant difference in the count of *U. affinis* in vials with and without *Aprostocetus* sp. emergence (*t* = 0.672, *df* = 88). In addition to *Aprostocetus* sp. and *U. affinis*, three *P. cardui* females and 10 *U. quadrifasciata* (6 males and 4 females) emerged.

Seed heads not placed into rearing were dissected and inspected for *U. affinis* and *U. quadrifasciata* galls. Galls produced by these two *Urophora* species are structurally different, with *U. affinis* producing woody, lignified galls and

<sup>1</sup>Michigan Technological University, Cooperative Emerald Ash Borer Project, 5936 Ford Ct. Suite 200, Brighton, MI, 48116. (e-mail: jmmarsha@mtu.edu).

*U. quadrifasciata* producing papery, non-lignified galls (White and Korneyev 1989, Burkhardt and Zwölfer 2002). Only galls produced by *U. affinis* were encountered. A sample of 15 *U. affinis* galls was dissected and 8 contained multiple *Aprostocetus* sp., with a mean parasitism rate of  $8.36 \pm 4.07$  *Aprostocetus* sp. individuals per *U. affinis* gall.

The gregarious endoparasitism of *Aprostocetus* sp. on *U. affinis* caused mortality in *U. affinis* at the sites in Livingston County, MI, where seed heads were collected. While this mortality did reduce the mean number of *U. affinis* individuals emerging within vials, it did not create significant differences in the vials with and without *Aprostocetus* sp. and did not significantly influence the presence of *U. affinis* individuals. With this first record of *Aprostocetus* sp. parasitizing *U. affinis*, further investigations into host selection and the geographic distribution of this parasitoid within the introduced ranges of *Urophora* spp. are warranted.

### ACKNOWLEDGMENTS

The author would like to thank Roger A. Burks (University of California, Riverside) for the identification of *Aprostocetus* sp. and Jonathan P. Lelito (The Pennsylvania State University) for lab assistance. Voucher specimens are held in the personal collection of the author.

### LITERATURE CITED

- Burkhardt, B., and H. Zwölfer. 2002. Macro-evolutionary tradeoffs in the tephritid *Urophora*: benefits and costs of an improved plant gall. *Evol. Ecol. Research* 4: 61-77.
- Foote, R. H., F. L. Blanc, and A. L. Norrbom. 1993. Handbook of the Fruit Flies (Diptera: Tephritidae) of America North of Mexico. Cornell University Press, Ithaca, New York.
- Harris, P. 1980. Establishment of *Urophora affinis* Frfld. and *U. quadrifasciata* (Meig.) (Diptera: Tephritidae) in Canada for the biological control of diffuse and spotted knapweed. *Z. Angew. Entomol.* 89: 504-514.
- Graham, M. W. R. de V. 1987. A reclassification of the European Tetrastichinae (Hymenoptera: Eulophidae), with a revision of certain genera. *Bull. Brit. Mus. (Nat. Hist.)*, Entomol. Series 55:1-392.
- Lang, R. F., R. D. Richard, and R. W. Hansen. 1997. *Urophora affinis* and *U. quadrifasciata* (Diptera: Tephritidae) released and monitored by USDA, APHIS, PPQ as biological control agents of spotted and diffuse knapweed. *Gt. Lakes Entomol.* 30: 105-113.
- LaSalle, J. 1994. North American genera of Tetrastichinae (Hymenoptera: Eulophidae). *J. Nat. Hist.* 28: 109-236.
- Marshall, J. M., R. A. Burks, and A. J. Storer. 2004. First host record for *Pteromalus cardui* (Hymenoptera: Pteromalidae) on *Urophora quadrifasciata* (Diptera: Tephritidae) in spotted knapweed (*Centaurea biebersteinii*, Asteraceae) in Michigan, U.S.A. *Entomol. News* 115: 273-278.
- Mays, W. T., and L. T. Kok. 2003. Population dynamics and dispersal of two exotic biological control agents of spotted knapweed, *Urophora affinis* and *U. quadrifasciata* (Diptera: Tephritidae), in southwestern Virginia from 1986 and 2000. *Biological Control* 27: 43-52.
- Noyes, J. S. 2003. Universal Chalcidoidea Database. The Natural History Museum, London. Available at: <<http://www.nhm.ac.uk/entomology/chalcidoids/>>
- Pearson, D. E. 1999. Deer mouse predation on the biological control agent, *Urophora* spp., introduced to control spotted knapweed. *Northwest. Nat.* 80: 26-29.

- Story, J. M., K. W. Boggs, W. R. Good, L. J. White, and R. M. Nowierski. 1995. Cause and extent of predation on *Urophora* spp. larvae (Diptera: Tephritidae) in spotted knapweed capitula. *Environ. Entomol.* 24: 1467-1472.
- White, I. M., and V. A. Korneyev. 1989. A revision of the western Palaearctic species of *Urophora* Robineau-Desvoidy (Diptera: Tephritidae). *Systematic Entomol.* 14: 327-374.
- Yegorenkova, E. N., Z. A. Yefremova, and V. V. Kostjukov. 2007. Contributions to the knowledge of tetrastichine wasps (Hymenoptera, Eulophidae, Tetrastichinae) of the Middle Volga Region. *Entomol. Review* 87: 1180-1192.