

Spring 2012

Evidence for the Impact of Reactive Oxygen Species on Branch Density Homeostasis in *Neurospora crassa*

Jacob Yablonowski
Valparaiso University

Taylor Grashel
Valparaiso University

Alex Zapata
Valparaiso University

Hamzah Abduljabar
Valparaiso University

Michael Watters
Valparaiso University

Follow this and additional works at: <https://scholar.valpo.edu/cus>

Recommended Citation

Yablonowski, Jacob; Grashel, Taylor; Zapata, Alex; Abduljabar, Hamzah; and Watters, Michael, "Evidence for the Impact of Reactive Oxygen Species on Branch Density Homeostasis in *Neurospora crassa*" (2012). Celebration of Undergraduate Scholarship. Paper 169

This Poster Presentation is brought to you for free and open access by the Office of Sponsored and Undergraduate Research at ValpoScholar. It has been accepted for inclusion in Symposium on Undergraduate Research and Creative Expression (SOURCE) by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

Evidence for the Impact of Reactive Oxygen Species on Branch Density Homeostasis in *Neurospora crassa*

Jacob Yablonowski, Tayler Grashel, Alex Zapata, Hamzah Abduljabar, Michael Watters

Departmental Affiliation: Biology
College of Arts and Sciences

In preliminary screenings, genes involved in the control of reactive oxygen species (ROS), were identified as playing a role in the process of growth rate compensation of branch density. Here we examine the relationship between ROS and branch density further. The maintenance of branch density under growth at various temperatures was examined in a selection of mutants in genes known to be important in ROS control. In all ROS control mutants tested, growth showed clear sustained hypobranching when grown at lower temperatures. This can be contrasted with wild-type *Neurospora* which branches at the same density under both conditions. We also tested the impact of environmental agents which lower the concentration of ROS on branching. In tests on wild type *Neurospora*, water soluble anti-oxidants (reducing agents), Ascorbic Acid and Glutathione produced unusual branching patterns. While lower doses produce sustainable hypobranching in wild-type *Neurospora*, hypha exposed to higher concentrations of Ascorbic Acid or Glutathione display a distribution of branching with two clear maxima. They show an increase in both very closely spaced branching as well as an increase in more distantly spaced branching.

Information about the Authors:
No information provided.

Faculty Sponsor: Dr. Michael Watters

Student Contact: jacob.yablonowski@valpo.edu