

April 1988

Ectoparasites and Other Arthropod Associates of the Hairy-tailed Mole, *Parascalops Breweri*

John O. Whitaker Jr
Indiana State University

Thomas W. French
Division of Fisheries and Wildlife

Follow this and additional works at: <http://scholar.valpo.edu/tgle>

 Part of the [Entomology Commons](#)

Recommended Citation

Whitaker, John O. Jr and French, Thomas W. (1988) "Ectoparasites and Other Arthropod Associates of the Hairy-tailed Mole, *Parascalops Breweri*," *The Great Lakes Entomologist*: Vol. 21 : No. 1 , Article 8.
Available at: <http://scholar.valpo.edu/tgle/vol21/iss1/8>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

ECTOPARASITES AND OTHER ARTHROPOD ASSOCIATES OF THE HAIRY-TAILED MOLE, *PARASCALOPS BREWERI*

John O. Whitaker, Jr.¹ and Thomas W. French²

ABSTRACT

A total of 33 taxa of ectoparasites and other associates was taken on seven individuals of the Hairy-tailed Mole, *Parascalops breweri*, from New York and New England. The most abundant form was the glycyphagid mite, *Labidophorus nearcticus*.

There are relatively few records of parasites, especially smaller mites, of the hairy-tailed mole, *Parascalops breweri* (Bachman). The summary of information on mites from this species by Whitaker and Wilson (1974) included only three species: a glycyphagid, *Labidophorus nearcticus* Fain and Whitaker (*Labidophorus talpae* as reported by Fain and Whitaker [1973]), and two laelapids, *Haemogamasus liponyssoides* Ewing reported by Keegan (1951) and *Echinonyssus blarinae* (Herrin) reported by Herrin (1970). Yates et al. (1979) examined ectoparasites from 23 hairy-tailed moles. These harbored a louse, *Haematopinoidea squamosus* (Osborn), a flea, *Ctenophthalmus pseudagyrtis* Baker, a beetle, *Leptinus testaceus* Mueller, and two laelapid mites, *Androlaelaps fahrenheitsi* (Berlese) and *Haemogamasus reidi* Ewing. *Pygmephorus whitakeri* Mahunka (Pygmephoridae) was reported from this host by Smiley and Whitaker (1979) and *Pygmephorus proctorae* Smiley and Whitaker was described from this host by Smiley and Whitaker (1984). Thus only 10 species of ectoparasites and other associates were previously known from this host.

We have examined seven hairy-tailed moles, four from the vicinity of Ithaca, Tompkins County, New York (taken 21 Sept.–8 Nov. 1980); one from DeBruce, Mongaup Creek, Ulster County, New York (11 Oct. 1981); one from Grafton County, New Hampshire 1.5 mi upstream from the Barnet Bridge of the Connecticut River (30 Aug. 1983); and one from Whitingham, Windham County, Vermont (4 Oct. 1983). The purpose of this paper is to present the results of this examination. Even though the host sample is small, the results are of interest, as we have identified from these moles 33 taxa of ectoparasites and other associates including at least 16 taxa of pygmephorids. Little is known of the biology and host relationships of pygmephorids. Only phoretomorphic females of *Pygmephorus* are found on mammalian hosts; these show little host specificity and are found most commonly on burrowers. Voucher specimens of many of the species are being deposited in the U.S. National Museum.

MATERIALS AND METHODS

Moles were collected in snap traps and immediately placed in plastic bags upon recovery. They were examined by manipulating their fur with dissecting needles while

¹Department of Life Sciences, Indiana State University, Terre Haute, IN 47809.

²Nongame and Endangered Species Program, Division of Fisheries and Wildlife, 100 Cambridge Street, Boston, MA 02202.

Table 1. Ectoparasites and other associates of seven hairy-tailed moles from New York and New England.

	Number		Frequency		Localities
	n	x	n	%	
Anoplura					
Hoplopleuridae					
<i>Haematopinoides squamosus</i>	3	0.43	1	14.3	NY
Siphonaptera					
Hystrichopsyllidae					
<i>Ctenophthalmus pseudagyrtis</i>	10	1.43	5	71.4	NY, NH
** <i>Doratopsylla blarinae</i> C. Fox	1	0.14	1	14.3	NY
** <i>Hystrichopsylla tahavuana</i>	3	0.43	2	28.6	NY
** <i>Nearctopsylla genalis</i> (Baker)	2	0.29	2	28.6	NH
Ceratophyllidae					
** <i>Megabothris asio</i> (Baker)	1	0.14	1	14.3	NH
Acari					
Histiostomatidae					
3	0.43	1	14.3	NY	
Ascidae					
<i>Proctolaelaps</i> sp.	6	0.86	4	57.1	NY, NH
Ologamasidae					
<i>Cyrtolaelaps</i> sp.	4	0.57	3	42.9	NY
<i>Euryparasitus</i> sp.	2	0.29	2	28.6	NY
Glycyphagidae					
<i>Labidophorus nearcticus</i>	276	39.43	5	71.4	NY, *NH, *VT
Laelapidae					
<i>Echinonyssus blarinae</i>	3	0.43	2	28.6	NY
<i>Haemogamasus liponyssoides</i>	4	0.57	1	14.3	*NH
** <i>Hypoaspis miles</i>	1	0.14	1	14.3	*NH
** <i>Laelaps alaskensis</i>	1	0.14	1	14.3	*NH
Pygmephoridae					
** <i>Bakerdania plurisetosa</i>	104	14.9	3	42.9	*NY, *VT
<i>Bakerdania</i> sp.	34	4.86	3	42.9	NY, VT
** <i>Pygmephorus brevicaudae</i> Smiley and Whitaker	1	0.14	1	14.3	*NY
** <i>P. designatus</i> Mahunka	1	0.14	1	14.3	*NY
** <i>P. hamiltoni</i> Smiley and Whitaker	4	0.57	3	42.9	*NY
** <i>P. hastatus</i> Mahunka	6	0.86	3	42.9	*NY
** <i>P. horridus</i> Mahunka	1	0.14	1	14.3	*NY
<i>P. proctorae</i>	28	4.00	4	57.1	NY, *NH
** <i>P. johnstoni</i> Smiley and Whitaker	2	0.29	1	14.3	*NY, *VT
** <i>P. lutterloughae</i> Smiley and Whitaker	2	0.29	1	14.3	*NY
** <i>P. rackae</i> Smiley and Whitaker	1	0.14	1	14.3	*NY
** <i>P. spinosus</i> Kramer	1	0.14	1	14.3	*NY
** <i>P. tamiasi</i> Mahunka	2	0.29	1	14.3	*NY, *NH, *VT
** <i>P. whartoni</i> Smiley and Whitaker	1	0.14	1	14.3	*NY
<i>P. whitakeri</i>	62	8.86	7	100.0	*NY, *NH, *VT
** <i>P. wrenschae</i> Smiley and Whitaker	1	0.14	1	14.3	*NY
<i>Pygmephorus</i> sp.	1	0.14	1	14.3*	NY
Trombiculidae					
<i>Neotrombicula microti</i> (Ewing)	5	0.71	1	14.3	VT

*New state records

**New host records

using a binocular microscope. Parasites were preserved in 70% alcohol, cleared and stained in Nesbitt's solution, mounted in Hoyer's solution, and the cover slips ringed with Euparal.

RESULTS AND DISCUSSION

The results are given in Table 1. A total of 33 taxa of ectoparasites and associates was taken from the seven moles, with the mites, *Labidophorus nearcticus*, *Bakerdania* spp. including *B. plurisetosa* Mahunka, *Pygmephorus whitakeri*, and *P. proctorae* being the most abundant. Pygmephorids, although not parasitic, were abundant, with a total of 252 individuals. Several species of pygmephorids were often taken on a single mole. This is many more species, 16, than has previously been taken on any one species of host. Four species of laelapid mites were reported earlier, and four were reported during this study, with *Laelaps alaskensis* Grant and *Hypoaspis miles* (Berlese) being new parasites from this host. However, only one individual of each was taken and both must be considered as accidentals.

Three individuals of the louse *Haematopinoides squamosus* were found. This species is rather uncommon occurring only on moles. Only one species of flea had been previously reported on this host, but we took five, including three individuals of the relatively rare species, *Hystrichopsylla tahaviana* Jordan.

New state and new host records are indicated by one or two asterisks, respectively, in Table 1.

LITERATURE CITED

- Fain, A., and J. O. Whitaker, Jr. 1973. Phoretic hypopi of North American mammals (Acarina: Sarcoptiformes, Glycyphagidae). *Acarologia* 15:144-170.
- . 1985. *Labidophorus nearcticus* n. sp. (Astigmata: Labidophorinae) a new glycyphagid mite from *Parascalops breweri* in the USA. *J. Parasitol.* 71:327-330.
- Herrin, C. S. 1970. A systematic revision of the genus *Hirstionyssus* (Acari: Mesostigmata) of the nearctic region. *J. Med. Entomol.* 7:391-437.
- Keegan, H. L. 1951. The mites of the subfamily Haemogamasinae (Acari: Laelaptidae). *Proc. U.S. Nat. Mus.* 101:203-268.
- Smiley, R. L., and J. O. Whitaker, Jr. 1979. Mites of the genus *Pygmephorus* (Acari: Pygmephoridae) on small mammals in North America. *Acta Zool. Acad. Scient. Hungarica.* 25:383-408.
- Whitaker, J. O., Jr., and N. Wilson. 1974. Host and distribution lists of mites (Acari), parasitic and phoretic, in the hair of wild mammals of North America, north of Mexico. *Amer. Midl. Nat.* 91:1-67.
- Yates, T. L., D. B. Pence, and G. K. Launchbaugh. 1979. Ectoparasites from seven species of North American moles (Insectivora: Talpidae). *J. Med. Entomol.* 16:166-168.